

Anti-Human IgG F(c) Secondary Antibody

Rabbit Polyclonal, Unconjugated Catalog # ASR3531

Specification

Anti-Human IgG F(c) Secondary Antibody - Product Information

Description Anti-HUMAN IgG F(c) (RABBIT) Antibody

Host Rabbit

Conjugate
Target Species
Clonality
Application

Unconjugated
Human
Polyclonal
WB, IHC, E

Application Note ELISA 1:20,000-1:100,000; Western Blot

1:2,000-1:10,000;Immunohistochemistry

1:1,000-1:5,000 Lyophilized

Physical State
Host Isotype
Target Isotype
Lyophilize
IgG
IgG F(c)

Buffer 0.01 M Sodium Phosphate, 0.15 M Sodium

Chloride, pH 7.2

Immunogen Human IgG F(c) fragment

Reconstitution Volume 5.0 n

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer None Preservative None

Anti-Human IgG F(c) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Human IgG, Human IgG F(c) and Human Serum. No reaction was observed against Human IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Human IgG F(c) Secondary Antibody - Protein Information



Anti-Human IgG F(c) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Human IgG F(c) Secondary Antibody - Images