

## Anti-Rabbit IgG F(c) Secondary Antibody

Donkey Polyclonal, Unconjugated Catalog # ASR3540

## **Specification**

Physical State Host Isotype

Target Isotype

**Immunogen** 

## Anti-Rabbit IgG F(c) Secondary Antibody - Product Information

Description Anti-RABBIT IgG F(c) (DONKEY) Antibody

Host Donkey
Conjugate Unconjugated
Target Species Rabbit
Clonality Polyclonal
Application ,1,10,15,

Application Note ELISA 1:20,000-1:100,000; Western Blot

1:2,000-1:10,000;Immunochemistry

1:1,000-1:5,000 Lyophilized

> IgG IgG F(c)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2
Rabbit IgG F(c) fragment

5.0 mL

Restore with deionized water (or

equivalent)

Preservative 0.01% (w/v) Sodium Azide

#### Anti-Rabbit IgG F(c) Secondary Antibody - Additional Information

## **Shipping Condition**

Reconstitution Volume Reconstitution Buffer

**Ambient** 

#### **Purity**

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Rabbit IgG, Rabbit IgG F(c) and Rabbit Serum. No reaction was observed against Rabbit IgG F(ab')2.

#### **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

## **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## Anti-Rabbit IgG F(c) Secondary Antibody - Protein Information



# **Anti-Rabbit IgG F(c) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Rabbit IgG F(c) Secondary Antibody - Images