

**Anti-HICE1 (Rabbit) Antibody**  
**HICE1 Antibody**  
**Catalog # ASR3877**

**Specification**

---

**Anti-HICE1 (Rabbit) Antibody - Product Information**

Host	Rabbit
Conjugate	Unconjugated
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	WB, E, I, LCI
Application Note	This antiserum has been tested for use in ELISA and western blotting using a full length recombinant Hice1 protein. Specific conditions for reactivity and detection of Hice1 should be optimized by the end user. Expect a band approximately ~45 kDa in size corresponding to Hice1 by Western Blotting in the appropriate cell lysate or extract.
Physical State	Liquid (sterile filtered)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Anti-HICE1 was prepared from whole rabbit serum produced by repeated immunizations with a recombinant full length Hice1 protein.
Preservative	0.01% (w/v) Sodium Azide

**Anti-HICE1 (Rabbit) Antibody - Additional Information**

**Gene ID** 93323

**Other Names**  
93323

**Purity**

This product was adsorbed against GST from monospecific antiserum by immunoaffinity chromatography. This antibody reacts with endogenous Hice1 protein. A BLAST analysis was used to suggest reactivity with Hice1 from human based on a 100% homology with the immunizing sequence. Expect reactivity with Hice1 from chimpanzee, Sumatran orangutan based on a 90% homology with the immunizing sequence. Cross-reactivity with Hice1 from other sources has not been determined.

**Storage Condition**

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

### Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

### Anti-HICE1 (Rabbit) Antibody - Protein Information

**Name** HAUS8

**Synonyms** HICE1

#### Function

Contributes to mitotic spindle assembly, maintenance of centrosome integrity and completion of cytokinesis as part of the HAUS augmin-like complex.

#### Cellular Location


Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, spindle pole. Note=During interphase, primarily cytoplasmic and associates with centrosomes and with the mitotic spindles, preferentially at the spindle pole vicinity. During anaphase and telophase, additionally associates with the spindle midzone and midbody, respectively. Localizes to mitotic spindle microtubules

### Anti-HICE1 (Rabbit) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-HICE1 (Rabbit) Antibody - Images



Anti-HICE1 in Western Blot using Rockland Immunochemicals' Anti-HICE1 Antibody shows detection of a 45 kDa band corresponding to endogenous HICE1 in lysates of S phase HeLa cells silenced for either control Luciferase or HICE1. In right lane (HICE1i): lysates from sh-HICE1 RNAi-treated lentivirus-infected cells. In left lane (GLi): lysates from sh-Luciferase

lentivirus-infected cells as control. Anti-HICE1 Antibody was used at 1:10,000. Molecular weight estimation was made by comparison by prestained MW markers. ECL was used for detection. Personal communication, Kyung S. Lee, NCI, Bethesda, MD.

### **Anti-HICE1 (Rabbit) Antibody - Background**

This antibody is designed, produced, and validated as part of a collaboration between Rockland and the National Cancer Institute (NCI) and is suitable for Cancer, Immunology and Nuclear Signaling research. Hice1 contributes to the mitotic spindle assembly, maintenance of centrosome integrity and completion of cytokinesis as part of the HAUS augmin-like complex. Normal bipolar spindle formation is critical for accurate chromosome segregation and proper mitotic progression. Failure in this event leads to spindle checkpoint activation and chromosome missegregation that ultimately leads to aneuploidy. Hice1 binds to microtubules directly, and promotes spindle integrity and chromosome stability. Hice1 has also shown to play an important role in targeting the gamma TuRC complex to the mitotic spindle, a step that appears to be required for spindle-mediated microtubule generation and normal chromosome segregation. The HAUS augmin-like complex's interaction with microtubules is strong during mitosis, while it is weak or absent during interphase. During interphase, it is primarily cytoplasmic, associating with centrosomes and with the mitotic spindles, preferentially at the spindle pole vicinity. During anaphase and telophase, it additionally associates with the spindle midzone and midbody, respectively. Further characterization of the function of Hice1 will likely be important for better understanding the mechanism of normal mitotic progression and high fidelity chromosome segregation.