

**Anti-Histone H3 (Rabbit) Antibody**  
**Histone H3 Antibody**  
**Catalog # ASR4467****Specification****Anti-Histone H3 (Rabbit) Antibody - Product Information**

Host	<b>Rabbit</b>
Conjugate	<b>Unconjugated</b>
Target Species	<b>Human</b>
Reactivity	<b>Human</b>
Clonality	<b>Polyclonal</b>
Application	<b>WB, IHC, E, I, LCI</b>
Application Note	<b>Histone H3 antibody has been tested for use in ELISA and western blot. This antibody is suitable for IHC. For western blots expect a band of approximately 15.4 kDa in size corresponding to the Histone 3 protein. Specific conditions for reactivity should be optimized by the end user.</b>
Physical State	<b>Liquid (sterile filtered)</b>
Buffer	<b>0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Immunogen	<b>Anti-Histone-3 was prepared from whole rabbit serum produced by repeated immunizations with a peptide corresponding to the c-terminus region of human histone-3.</b>
Preservative	<b>0.01% (w/v) Sodium Azide</b>

**Anti-Histone H3 (Rabbit) Antibody - Additional Information****Gene ID** 8350;8351;8352;8353;8354;8355;8356;8357;8358;8968**Other Names**

8350

**Purity**

Anti-Histone H3 is directed against the human histone 3 protein. The product was protein A purified from monospecific antiserum by immunoaffinity chromatography using protein A coupled to agarose beads. A BLAST analysis was used to suggest reactivity with human and multiple other eukaryotic. Cross-reactivity with histone-3 from other sources have not been determined.

**Storage Condition**

Store H3 Antibody at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## Anti-Histone H3 (Rabbit) Antibody - Protein Information

**Name** H3C1 ([HGNC:4766](#))

**Synonyms** H3FA, HIST1H3A

### Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

### Cellular Location


Nucleus. Chromosome.

## Anti-Histone H3 (Rabbit) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-Histone H3 (Rabbit) Antibody - Images



Western Blot of Rabbit Anti-Histone H3 antibody. Lane 1: HeLa WCL (p/n W09-000-364). Lane 2: HeLa Nuclear Extract (p/n W09-001-367). Load: 10 µg per lane. Primary antibody: Histone H3 antibody at 1.0 µg/ml for 1 hr at RT. Secondary antibody: HRP Gt-a-Rb IgG secondary antibody (p/n 611-103-122) at 1:40,000 for 30 min at RT. Block: MB-070 overnight at 4°C. Predicted/Observed size: ~15 kDa, ~15 kDa for Histone H3.

## Anti-Histone H3 (Rabbit) Antibody - Background

Histone H3 is one of the five main histone proteins involved in the structure of chromatin in eukaryotic cells. Histone proteins are highly post-translationally modified with Histone H3 being the most extensively modified of the five histones. The N-terminal tail of histone H3 protrudes from the globular nucleosome core and can undergo several different types of post-translational modification that influence cellular processes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Histone H3 Antibody is ideal for investigators involved in Cell Signaling, Epigenetics, Nuclear Signaling research and Signal Transduction research.