

Anti-LACTOPEROXIDASE (Bovine Milk) (SHEEP) Antibody Biotin
Anti-Lactoperoxidase Antibody Biotin Conjugated
Catalog # ASR4827**Specification****Anti-LACTOPEROXIDASE (Bovine Milk) (SHEEP) Antibody Biotin - Product Information**

Host	Sheep
Conjugate	Biotin
Target Species	Bovine
Reactivity	Bovine
Clonality	Polyclonal
Application	WB, E, IP, I, LCI
Application Note	Anti-Lactoperoxidase Biotin Conjugated Antibody has been tested by Dot blot and western blot and is suitable to be assayed against 1.0 ug of Lactoperoxidase in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:1,000 to 1:6,000 of the reconstitution concentration is suggested for Anti-Lactoperoxidase.
Physical State	Lyophilized
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Lactoperoxidase [Bovine Milk]
Reconstitution Volume	100 µL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

Anti-LACTOPEROXIDASE (Bovine Milk) (SHEEP) Antibody Biotin - Additional Information**Gene ID** 280844**Other Names**
280844**Purity**

Anti-Lactoperoxidase (Bovine Milk) (Sheep) Antibody Biotin Conjugated is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Sheep Serum as well as purified and partially purified Lactoperoxidase [Bovine Milk]. Cross reactivity against Lactoperoxidase from other sources is unknown.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-LACTOPEROXIDASE (Bovine Milk) (SHEEP) Antibody Biotin - Protein Information**Name** LPO**Function**

Heme-containing oxidoreductase which catalyzes the conversion of thiocyanate (SCN(-)) into antimicrobial agent hypothiocyanous acid (OSCN(-)) in the presence of hydrogen peroxide (H₂O₂) (Probable) (PubMed:5338806). Also involved in the conversion of iodide (I(-)) into hypoiodite (IO(-)) in the presence of H₂O₂ (Probable) (PubMed:354515). Responsible for the inactivation of a wide range of micro-organisms and hence, important component of defense mechanism (PubMed:354515, PubMed:5338806). The lactoperoxidase-SCN(-)-H₂O₂ system shows antibacterial properties against some streptococci strains (PubMed:5338806). The lactoperoxidase-I(-)-H₂O₂ system shows antibacterial properties against E.coli (PubMed:354515). May protect the udder from infection and may promote growth in newborns (By similarity). May be implicated in airway host defense against infection (By similarity). May contribute to maintaining an appropriate H₂O₂ cellular level, therefore protecting cells from H₂O₂-caused injuries and inflammation (By similarity).

Cellular Location

Secreted. Cytoplasm {ECO:0000250|UniProtKB:Q5SW46}

Tissue Location

Mammary gland; milk..

Anti-LACTOPEROXIDASE (Bovine Milk) (SHEEP) Antibody Biotin - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-LACTOPEROXIDASE (Bovine Milk) (SHEEP) Antibody Biotin - Images

Anti-LACTOPEROXIDASE (Bovine Milk) (SHEEP) Antibody Biotin - Background

Lactoperoxidase antibody recognizes the lactoperoxidase protein. Lactoperoxidase catalyzes the oxidation of a number of inorganic and organic substrates by hydrogen peroxide. Lactoperoxidase plays an important role in killing bacteria in milk. Lactoperoxidase conjugated to Biotin is used to detect the specific target of the lactoperoxidase protein. Lactoperoxidase conjugated to Biotin is suitable for researchers in immunology and biochemistry.