

Anti-PC2 [RABBIT] Antibody
PC2 Antibody
Catalog # ASR5251**Specification**

Anti-PC2 [RABBIT] Antibody - Product Information

Host	Rabbit
Conjugate	Unconjugated
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	WB, E, IP, I, LCI
Application Note	Anti-PC2 antibody has been tested for use in ELISA, WB, and IF. This affinity purified antibody is useful in immunofluorescence staining of cultured cells. In immunofluorescence, this antibody detects the expected discrete nuclear structure that is termed the PcG body, corresponding to the known localization of PC2. This affinity purified antibody is useful in western blotting using transfected cell lysates. Dilutions for western blotting represent a starting point dilution and further optimization may be required. The antibody detects a band of approximately 82 kDa (predicted molecular weight: 61.4 kDa). The antibody has been successfully used to detect FLAG-tagged transfected hPC2 (see figure). It detects a weak band that probably corresponds to endogenous PC2, however, a strong secondary band is also seen at 50kD in all cell lines thus far tested. This suggests that the antibody may also react with another highly expressed ubiquitous protein. Reactivity in other immunoassays is unknown.
Physical State	Liquid (sterile filtered)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 90-115 of Human PC2 protein.
Preservative	0.01% (w/v) Sodium Azide

Anti-PC2 [RABBIT] Antibody - Additional Information

Gene ID 8535**Other Names**
8535**Purity**

This is an affinity purified antibody produced by immunoaffinity chromatography using the immunizing peptide after immobilization to a solid phase. Reactivity occurs against human PC2 protein.

Storage Condition

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-PC2 [RABBIT] Antibody - Protein Information**Name** CBX4**Function**

E3 SUMO-protein ligase that catalyzes sumoylation of target proteins by promoting the transfer of SUMO from the E2 enzyme to the substrate (PubMed:12679040, PubMed:22825850). Involved in the sumoylation of HNRNPK, a p53/TP53 transcriptional coactivator, hence indirectly regulates p53/TP53 transcriptional activation resulting in p21/CDKN1A expression. Monosumoylates ZNF131 (PubMed:22825850).

Cellular Location

Nucleus. Nucleus speckle. Note=Localization to nuclear polycomb bodies is required for ZNF131 sumoylation (PubMed:22467880). Localized in distinct foci on chromatin (PubMed:18927235)

Tissue Location


Ubiquitous.

Anti-PC2 [RABBIT] Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PC2 [RABBIT] Antibody - Images



Rockland's anti-hPC2 antibody was used for immunofluorescent imaging of human cells (U2OS). The image reveals the expected discrete nuclear structure that is termed the PcG body corresponding to the known localization of PC2 (see Satijn et al. below). IF was performed after fixation in PBS with 4% PF for 5 min, permeabilization with 0.5% Triton X100-PBS for 5 min, and blocking with 5% milk / 0.2% Tween for 1 h. Primary antibody used at 1:200 in 5% milk / 0.2% Tween for 1 h, secondary antibody for 30 min. All blocking and incubation steps carried out at 37° C. Nuclei were counterstained with Hoechst stain (blue). Data contributed by Luke Hughes-Davies and Rhiannon Jade, Gurdon Institute, Cambridge, UK.

Anti-PC2 [RABBIT] Antibody - Background

PC2 is the human homolog of the *Drosophila* 'Polycomb' (Pc) protein which has been identified as a gene family member associated with a cellular memory system that is responsible for the inheritance of gene activity by progeny cells. The human Pc homolog gene is more closely related to a *Xenopus* Pc homolog, XPc, than to a previously described human Pc homolog, CBX2 (hPc1). However, the hPc2 and CBX2/hPc1 proteins are shown to colocalize in interphase nuclei of human U-2 OS osteosarcoma cells, suggesting that the proteins are part of a common protein complex. The human protein is believed to function as a repressor of proto-oncogene activity and that interference with hPc2 function can lead to depression of proto-oncogene transcription and subsequently to cellular transformation. Other reports describe PC2 as a protein that has SUMO E3 activity for the corepressors CtBP and CtBP2.