

**Anti-SUMO Activating Enzyme E1 (SAE1) pS185 (RABBIT) Antibody**  
**SAE1 phospho S185 Antibody**  
**Catalog # ASR5471****Specification****Anti-SUMO Activating Enzyme E1 (SAE1) pS185 (RABBIT) Antibody - Product Information**

Host	Rabbit
Conjugate	Unconjugated
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	WB, E, I, LCI
Application Note	This purified antibody has been tested for use in ELISA and western blot. Specific conditions for reactivity should be optimized by the end user. Expect a band at ~37 kDa in size corresponding to phosphorylated SAE1 protein by western blotting in the appropriate cell lysate or extract. This phospho-specific antibody reacts with human SAE1 pS185 and shows minimal reactivity by ELISA against the non-phosphorylated form of the immunizing peptide.
Physical State	Liquid (sterile filtered)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	This purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to a region surrounding S185 of the human SUMO Activating Enzyme E1 protein.
Preservative	0.01% (w/v) Sodium Azide

**Anti-SUMO Activating Enzyme E1 (SAE1) pS185 (RABBIT) Antibody - Additional Information****Gene ID** 10055**Other Names**  
10055**Purity**

This purified antibody is directed against human SUMO Activating Enzyme E1 protein. The product was purified from monospecific antiserum by affinity chromatography. This antibody is specific for human SAE1 protein phosphorylated at S185. A BLAST analysis using the sequence of the immunizing peptide was used to suggest that this antibody would react with SUMO Activating Enzyme E1 protein from human (100%), bovine, dog, chimpanzee (96%), mouse (93%), and rat (92%) based on a high degree of sequence homology. Cross reactivity against this protein from

other sources has not been determined.

**Storage Condition**

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Anti-SUMO Activating Enzyme E1 (SAE1) pS185 (RABBIT) Antibody - Protein Information**

**Name** SAE1

**Synonyms** AOS1, SUA1, UBLE1A

**Function**

The heterodimer acts as an E1 ligase for SUMO1, SUMO2, SUMO3, and probably SUMO4. It mediates ATP-dependent activation of SUMO proteins followed by formation of a thioester bond between a SUMO protein and a conserved active site cysteine residue on UBA2/SAE2.

**Cellular Location**

Nucleus.


**Tissue Location**


Expression level increases during S phase and drops in G2 phase (at protein level).

**Anti-SUMO Activating Enzyme E1 (SAE1) pS185 (RABBIT) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-SUMO Activating Enzyme E1 (SAE1) pS185 (RABBIT) Antibody - Images**



Western blot using Rockland's Rabbit anti-SAE1 pS185 antibody shows detection of phosphorylated SAE1. Left lane (-) contains 20 µg human HeLa whole cell protein. Right lane (+) contains 20 µg human HeLa whole cell protein from cells pre-treated with phosphatase inhibitor cocktail to prevent dephosphorylation of the target. Proteins were separated on a 10% SDS-PAGE and transferred onto nitrocellulose. After blocking with 5% milk-TBST 1 hr at room temperature, the membrane was probed with the primary antibody diluted to 1:1,000 at room temperature for 3 hr followed by washes and reaction with HRP-conjugated secondary and ECL imaging. Personal communication, Xin-Hua Feng, Baylor College of Medicine, Houston, TX

#### **Anti-SUMO Activating Enzyme E1 (SAE1) pS185 (RABBIT) Antibody - Background**

SUMO E1 activating enzyme (also called Ubiquitin-like 1 activating enzyme E1A, UBLE1A, AOS1, SAE1, and SUA1) with SAE2 (also known as UBA2) forms a heterodimeric (SAE1/SAE2) enzyme that activates the ubiquitin-like SUMO proteins (SUMO stands for Small Ubiquitin-like Modifier.) The SAE1 (SUMO Activating Enzyme 1) subunit resembles the N-terminal half of yeast UBA1; the SAE2 (also called Uba2) subunit corresponds to the C-terminal part of yeast UBA1 and contains the active site cysteine. In the SUMO activation step, SAE1/SAE2 uses ATP to adenylate the C-terminal glycine of SUMO-1 (the first of the three different mammalian SUMO proteins) then forms a high-energy thioester bond between the C-terminal glycine and the active site cysteine in SAE2 (Uba2). In the conjugation step, the SUMO moiety is transferred from SAE1/SAE2 to the active site cysteine (Cys 93) of the SUMO conjugating enzyme (SUMO E2, Ubc9) forming a SUMO-E2 thioester complex.