

Anti-GAB1 pY659 (RABBIT) Antibody

GAB1 phospho Y659 Antibody Catalog # ASR5523

Specification

Application Note

Physical State

Buffer

Anti-GAB1 pY659 (RABBIT) Antibody - Product Information

Host Rabbit

Conjugate
Target Species
Reactivity
Clonality
Application

Unconjugated
Human
Human
Polyclonal
WB, E, I, LCI

Anti-GAB1 pY656 Antibody has been tested for use in ELISA and by western blot.

Specific conditions for reactivity should be optimized by the end user. Expect a

predominant band at 76.6 kDa by western blotting in the appropriate cell lysate or extract. This phospho-specific polyclonal antibody reacts with human GAB1 pY659 and shows minimal reactivity by ELISA against the non-phosphorylated form of

the immunizing peptide. Liquid (sterile filtered)

0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen Anti-GAB1 pY656 Antibody was produced

in rabbits by repeated immunizations with

a synthetic peptide corresponding to

residues surrounding Y659 of human GAB1

protein.

Preservative 0.01% (w/v) Sodium Azide

Anti-GAB1 pY659 (RABBIT) Antibody - Additional Information

Gene ID 2549

Other Names

2549

Purity

Anti-GAB1pY659 was prepared from monospecific antiserum by immunoaffinity chromatography using phospho peptide coupled to agarose beads followed by solid phase adsorptions against non-phospho peptide and non-specific peptide to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum. Antibody is specific for human GAB1 protein phosphorylated at Y659. Cross-reactivity against GAB1 from other species may occur but has not yet been tested.

Storage Condition

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended



storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-GAB1 pY659 (RABBIT) Antibody - Protein Information

Name GAB1

Function


Adapter protein that plays a role in intracellular signaling cascades triggered by activated receptor-type kinases. Plays a role in FGFR1 signaling. Probably involved in signaling by the epidermal growth factor receptor (EGFR) and the insulin receptor (INSR). Involved in the MET/HGF-signaling pathway (PubMed:29408807).

Anti-GAB1 pY659 (RABBIT) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-GAB1 pY659 (RABBIT) Antibody - Images



Western Blot of Rabbit anti-GAB1 pY659 antibody. Load: Phospho GAB1 rProtein 0.05 μ g per lane. Blot Lane 1: GAB1 pY659 primary antibody at 1 μ g/mL in absence of blocking peptide overnight at 4°C. Blot Lane 2: GAB1 pY659 primary antibody at 1 μ g/mL incubated with 50 μ g of GAB1 pY659 peptide overnight at 4°C. Secondary antibody: HRP Goat Anti-Rabbit IgG secondary antibody (p/n 611-103-122) at 1:40,000 for 30 min at RT. Block: (p/n MB-070) Fluorescent blocking buffer for 30 min at RT. Predicted/Observed size: 130 kDa, 130 kDa for GAB1pY659. Other band(s): none.







Anti-GAB1 pY659 (RABBIT) Antibody - Background

GAB1 pY659 antibody detects phosphorylated GAB1. GAB1 is a member of the IRS1-like multisubstrate docking protein family. The protein is an important mediator of branching tubulogenesis and plays a central role in cellular growth response, transformation and apoptosis. Two transcript variants encoding different isoforms have been found for this gene. GAB1 plays a role in intracellular signaling cascades triggered by activated receptor-type kinases. It is known to play a role in FGFR1 signaling and is probably involved in signaling by the epidermal growth factor receptor (EGFR) and the insulin receptor (INSR). GAB1 interacts with GRB2 and with other SH2-containing proteins. It is known to interact with phosphorylated LAT2, PTPRI, FRS2, GRB2, PIK3R1 and SOS1. GAB1 gets phosphorylated in response to FGFR1 activation. This tyrosine phosphorylation of GAB1 mediates interaction with several proteins that contain SH2 domains. Anti-GAB1 Antibody is ideal for investigators involved in Cell Signaling, Cancer, Neuroscience and Signal Transduction research.