

Anti-Histone H3 [p Ser28] (RABBIT) Antibody
Histone H3 phospho S28 Antibody
Catalog # ASR5642**Specification**

Anti-Histone H3 [p Ser28] (RABBIT) Antibody - Product Information

Host	Rabbit
Conjugate	Unconjugated
Target Species	Human
Reactivity	Human, Mouse
Clonality	Polyclonal
Application	WB, IHC, I, LCI
Application Note	Anti-Histone H3 [p Ser28] antibody is tested for Western Blot and Immunofluorescence. This antibody is useful for Chromatin Immunoprecipitation, Dot Blot, and Immunocytochemistry. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately ~15.4 kDa corresponding to Histone H3 protein by Western Blotting in HeLa histone prep lysate or the appropriate cell lysate or extract. Epi-Plus™ antibody production in collaboration with Novus Biologicals.
Physical State	Liquid (sterile filtered)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Histone H3 [p Ser28] affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic phosphorylated peptide surrounding Serine 28 of human Histone H3.
Stabilizer	30% Glycerol
Preservative	0.01% (w/v) Sodium Azide

Anti-Histone H3 [p Ser28] (RABBIT) Antibody - Additional Information**Gene ID** 126961;333932;653604**Other Names**
126961**Purity**

Anti-Histone H3 [p Ser28] was affinity purified from monospecific antiserum by immunoaffinity chromatography. This antibody reacts with human Histone H3. A BLAST analysis was used to suggest cross-reactivity with Human, mouse, and C. elegans. Predicted to react with many species including rat, chicken, Xenopus, Drosophila, and plant based on 100% sequence homology. Cross-reactivity with Histone H3 from other sources has not been determined.

Storage Condition

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Histone H3 [p Ser28] (RABBIT) Antibody - Protein Information

Name H3C15 ([HGNC:20505](#))

Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.


Cellular Location

Nucleus. Chromosome.

Anti-Histone H3 [p Ser28] (RABBIT) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Histone H3 [p Ser28] (RABBIT) Antibody - Images

Immunofluorescence of Rabbit Anti-Histone H3 [p Ser28] Antibody. Tissue: HeLa cells. Fixation:

0.5% PFA. Antigen retrieval: Not required. Primary antibody: Histone H3 [p Ser28] antibody at a 1:500 dilution for 1 h at RT. Secondary antibody: Dylight 488 secondary antibody at 1:10,000 for 45 min at RT. Localization: Histone H3 [p Ser28] is nuclear and chromosomal. Staining: Histone H3 [p Ser28] is expressed in green, nuclei and alpha-tubulin are counterstained with DAPI (blue) and Dylight 550 (red).

Anti-Histone H3 [p Ser28] (RABBIT) Antibody - Background

Chromatin is the arrangement of DNA and proteins in which chromosomes are formed. Correspondingly, chromatin is formed from nucleosomes, which are comprised of a set of four histone proteins (H2A, H2B, H3, H4) wrapped with DNA. Chromatin is a very dynamic structure in which numerous post-translational modifications work together to activate or repress the availability of DNA to be copied, transcribed, or repaired. These marks decide which DNA will be open and commonly active (euchromatin) or tightly wound to prevent access and activation (heterochromatin). Common histone modifications include methylation of lysine and arginine, acetylation of lysine, phosphorylation of threonine and serine, and sumoylation, biotinylation, and ubiquitylation of lysine. In particular, phosphorylation of serine 28 of H3 (H3 pS28) is a mitotic and immediate early gene marker. Although similar to phosphorylation on serine 10 of H3 in the role of chromatin condensation in mitosis and the ability to respond to external stimuli such as stress and mitogen activation, these two phosphorylation sites are independent and promote gene expression separately. Phosphorylation of S28 is modulated by MSK1, MSK2, and Aurora-B kinase. Anti-Histone H3 are ideal for researchers interested in Chromatin Research, Epigenetics, Chromatin Modifiers, Histones and Modified Histones research.