

**Anti-HDAC9 (RABBIT) Antibody**  
**HDAC9 (N-terminus) Antibody**  
**Catalog # ASR5719****Specification**

---

**Anti-HDAC9 (RABBIT) Antibody - Product Information**

|                  |   |
|------------------|---|
| Host             | Rabbit  |
| Conjugate        | Unconjugated  |
| Target Species   | Human   |
| Reactivity       | Human   |
| Clonality        | Polyclonal  |
| Application      | WB, IHC, E, I, LCI  |
| Application Note | Anti-HDAC9 antibody is tested by Immunohistochemistry and useful for ELISA and Western Blot. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately ~111kDa corresponding to the appropriate cell lysate or extract. |
| Physical State   | Liquid (sterile filtered)   |
| Buffer           | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |
| Immunogen        | Anti-HDAC9 affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide at the N-terminal of human HDAC9 protein.  |
| Stabilizer       | 30% Glycerol  |

**Anti-HDAC9 (RABBIT) Antibody - Additional Information****Gene ID** 9734**Purity**

Anti-HDAC9 was affinity purified from monospecific antiserum by immunoaffinity chromatography. A BLAST analysis was used to suggest cross-reactivity with human based on 100% sequence homology. Cross-reactivity with HDAC9 from other sources has not been determined.

**Storage Condition**

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Anti-HDAC9 (RABBIT) Antibody - Protein Information**

**Name** HDAC9

**Synonyms** HDAC7, HDAC7B, HDRP, KIAA0744, MITR

**Function**

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Represses MEF2-dependent transcription.

**Cellular Location**

Nucleus.

**Tissue Location**


Broadly expressed, with highest levels in brain, heart, muscle and testis. Isoform 3 is present in human bladder carcinoma cells (at protein level).

**Anti-HDAC9 (RABBIT) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-HDAC9 (RABBIT) Antibody - Images**



Immunohistochemistry of Rabbit anti-HDAC9 antibody. Tissue: Small Intestine. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: HDAC9 antibody at 5 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Staining: HDAC9 as precipitated red signal with hematoxylin purple nuclear counterstain.

**Anti-HDAC9 (RABBIT) Antibody - Background**

HDAC9 is located in the nucleus, expressed most highly in brain, heart, muscle, and testis. It is

responsible for the deacetylation of lysine residues on the N-terminal region of the core histones (H2A, H2B, H3 and H4). The result of deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. HDAC9 represses MEF2-dependent transcription by recruiting HDAC1 and 3. It appears to inhibit skeletal myogenesis and be a factor in heart development. By repressing JUN transcription via HDAC1 and inhibiting JUN phosphorylation by MAPK10, HDAC9 protects neurons from apoptosis. Anti-HDAC9 therefore is ideal for investigators interested in Cardiovascular or Epigenetics and Nuclear Signaling research.