

Anti-Collagen Type I (RABBIT) Antibody Biotin Conjugated
Collagen Type I Antibody Biotin Conjugated
Catalog # ASR5849**Specification**

Anti-Collagen Type I (RABBIT) Antibody Biotin Conjugated - Product Information

| | |
|-----------------------|---|
| Host | Rabbit |
| Conjugate | Biotin |
| FP Value | 10-20 |
| Target Species | Mammalian |
| Reactivity | Rat, Human, Mouse |
| Clonality | Polyclonal |
| Application | WB, IHC, E, IP, I, LCI |
| Application Note | Anti-COLLAGEN Type I Antibody Biotin Conjugated has been tested by dot blot and Flow Cytometry and is suitable for western blot, immunoprecipitation, Flow Cytometry, and immunohistochemistry. Researchers should determine optimal titers for applications that are not stated below. |
| Physical State | Lyophilized |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Immunogen | Collagen Type I from human and bovine placenta. |
| Reconstitution Volume | 100 µL |
| Reconstitution Buffer | Restore with deionized water (or equivalent) |
| Stabilizer | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| Preservative | 0.01% (w/v) Sodium Azide |

Anti-Collagen Type I (RABBIT) Antibody Biotin Conjugated - Additional Information**Gene ID 1277****Other Names**
1277**Purity**

This product has been prepared by immunoaffinity chromatography using immobilized antigens. Some class-specific anti-collagens may be specific for three-dimensional epitopes which may result in diminished reactivity with denatured collagen or formalin-fixed, paraffin embedded tissues. This antibody reacts with most mammalian Type I collagens and has expected cross-reactivity with Type III and negligible cross reactivity with Type II, IV, V or VI collagens. Non-specific cross-reaction of anti-collagen antibodies with other human serum proteins or non-collagen extracellular matrix proteins has not been tested.

Storage Condition

Store vial at 4° C prior to restoration. Restore with 0.1 mL of deionized water (or equivalent). For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Collagen Type I (RABBIT) Antibody Biotin Conjugated - Protein Information

Name COL1A1

Function

Type I collagen is a member of group I collagen (fibrillar forming collagen).

Cellular Location

Secreted, extracellular space, extracellular matrix {ECO:0000255|PROSITE-ProRule:PRU00793}

Tissue Location


Forms the fibrils of tendon, ligaments and bones. In bones the fibrils are mineralized with calcium hydroxyapatite

Anti-Collagen Type I (RABBIT) Antibody Biotin Conjugated - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Collagen Type I (RABBIT) Antibody Biotin Conjugated - Images



Flow Cytometry of Anti-Collagen Type I Biotin Conjugated Antibody (600-406-103). Cells: mouse

lung. Stimulation: none. Primary antibody: biotin conjugated anti-collagen type I antibody. Secondary antibody: PE-conjugated CD45 and PE-conjugated anti-collagen type I secondary antibody.

Anti-Collagen Type I (RABBIT) Antibody Biotin Conjugated - Background

COLLAGEN Type I Antibody Biotin Conjugated is specific for Collagen Type I. Collagen Type I (Type-I collagen) is the most abundant collagen of the human body. It is present in scar tissue, the end product when tissue heals by repair. It is also found in tendons, the endomysium of myofibrils and the organic part of bone. Anti-collagen Type I antibody is suitable for Cancer research and other general research.