

ASAH1 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant ASAH1.****Catalog # AT1207a****Specification**

ASAH1 Antibody (monoclonal) (M01) - Product Information

Application	WB, IHC, E
Primary Accession	Q13510
Other Accession	NM_177924
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG3 Kappa
Calculated MW	44660

ASAH1 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 427**Other Names**

Acid ceramidase, AC, ACDase, Acid CDase, Acylsphingosine deacylase, N-acylsphingosine amidohydrolase, Putative 32 kDa heart protein, PHP32, Acid ceramidase subunit alpha, Acid ceramidase subunit beta, ASAH1, ASAH

Target/Specificity

ASAH1 (NP_808592, 25 a.a. ~ 124 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

IHC~~1:100~500

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions


ASAH1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

ASAH1 Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


ASAH1 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .



Immunoperoxidase of monoclonal antibody to ASAH1 on formalin-fixed paraffin-embedded human heart. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged ASAH1 is 0.1 ng/ml as a capture antibody.

ASAH1 Antibody (monoclonal) (M01) - Background

This gene encodes a heterodimeric protein consisting of a nonglycosylated alpha subunit and a glycosylated beta subunit that is cleaved to the mature enzyme posttranslationally. The encoded protein catalyzes the synthesis and degradation of ceramide into sphingosine and fatty acid. Mutations in this gene have been associated with a lysosomal storage disorder known as Farber disease. Multiple transcript variants encoding several distinct isoforms have been identified for this gene.

ASAH1 Antibody (monoclonal) (M01) - References

1.Ceramide biosynthesis and metabolism in trophoblast syncytialization.Singh AT, Dharmarajan A, Aye IL, Keelan JA.Mol Cell Endocrinol. 2012 May 28.