

ATP5G1 Antibody (monoclonal) (M01)
Mouse monoclonal antibody raised against a full length recombinant ATP5G1.
Catalog # AT1238a

Specification

ATP5G1 Antibody (monoclonal) (M01) - Product Information

Application	WB
Primary Accession	P05496
Other Accession	BC004963
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	14277

ATP5G1 Antibody (monoclonal) (M01) - Additional Information

Gene ID 516

Other Names

ATP synthase F(0) complex subunit C1, mitochondrial, ATP synthase lipid-binding protein, ATP synthase proteolipid P1, ATP synthase proton-transporting mitochondrial F(0) complex subunit C1, ATPase protein 9, ATPase subunit c, ATP5G1

Target/Specificity

ATP5G1 (AAH04963, 18 a.a. ~ 136 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

ATP5G1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


ATP5G1 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ATP5G1 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (38.83 KDa) .

ATP5G1 Antibody (monoclonal) (M01) - Background

This gene encodes a subunit of mitochondrial ATP synthase. Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. ATP synthase is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, comprising the proton channel. The catalytic portion of mitochondrial ATP synthase consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled with a stoichiometry of 3 alpha, 3 beta, and a single representative of the other 3. The proton channel seems to have nine subunits (a, b, c, d, e, f, g, F6 and 8). This gene is one of three genes that encode subunit c of the proton channel. Each of the three genes have distinct mitochondrial import sequences but encode the identical mature protein. Alternatively spliced transcript variants encoding the same protein have been identified.

ATP5G1 Antibody (monoclonal) (M01) - References

1. Neuronal cell surface ATP synthase mediates synthesis of extracellular ATP and regulation of intracellular pH. Xing SL, Yan J, Yu ZH, Zhu CQ. *Cell Biol Int.* 2010 Jul 14. [Epub ahead of print]