

**BOLL Antibody (monoclonal) (M09)****Mouse monoclonal antibody raised against a partial recombinant BOLL.****Catalog # AT1309a****Specification****BOLL Antibody (monoclonal) (M09) - Product Information**

Application	WB, E
Primary Accession	<a href="#">Q8N9W6</a>
Other Accession	<a href="#">NM_033030</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	31301

**BOLL Antibody (monoclonal) (M09) - Additional Information****Gene ID 66037****Other Names**

Protein boule-like, BOLL, BOULE

**Target/Specificity**

BOLL (NP\_149019, 185 a.a. ~ 283 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

E~~N/A

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**


BOLL Antibody (monoclonal) (M09) is for research use only and not for use in diagnostic or therapeutic procedures.

**BOLL Antibody (monoclonal) (M09) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)


- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**BOLL Antibody (monoclonal) (M09) - Images**

Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .



BOLL monoclonal antibody (M09), clone 1A7 Western Blot analysis of BOLL expression in Hela S3 NE ( (Cat # AT1309a )



Detection limit for recombinant GST tagged BOLL is approximately 0.3ng/ml as a capture antibody.

**BOLL Antibody (monoclonal) (M09) - Background**

This gene belongs to the DAZ gene family required for germ cell development. It encodes an RNA-binding protein which is more similar to *Drosophila* Boule than to human proteins encoded by genes DAZ (deleted in azoospermia) or DAZL (deleted in azoospermia-like). Loss of this gene function results in the absence of sperm in semen (azoospermia). Histological studies demonstrated that the primary defect is at the meiotic G2/M transition. Two alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.

#### **BOLL Antibody (monoclonal) (M09) - References**

Human DAZL, DAZ and BOULE genes modulate primordial germ-cell and haploid gamete formation. Kee K, et al. *Nature*, 2009 Nov 12. PMID 19865085. Posttranscriptional regulation of CDC25A by BOLL is a conserved fertility mechanism essential for human spermatogenesis. Lin YM, et al. *J Clin Endocrinol Metab*, 2009 Jul. PMID 19417033. Phenotypic expression of partial AZFc deletions is independent of the variations in DAZL and BOULE in a Han population. Chen P, et al. *J Androl*, 2010 Mar-Apr. PMID 19342699. Susceptibility loci for intracranial aneurysm in European and Japanese populations. Bilguvar K, et al. *Nat Genet*, 2008 Dec. PMID 18997786. Association of three isoforms of the meiotic BOULE gene with spermatogenic failure in infertile men. Kostova E, et al. *Mol Hum Reprod*, 2007 Feb. PMID 17114206.