

CABC1 Antibody (monoclonal) (M05A)

Mouse monoclonal antibody raised against a partial recombinant CABC1. Catalog # AT1362a

Specification

CABC1 Antibody (monoclonal) (M05A) - Product Information

Application WB **Primary Accession Q8NI60** Other Accession BC005171 Reactivity Human Host Mouse Clonality **Monoclonal** Isotype IgM Kappa Calculated MW 71950

CABC1 Antibody (monoclonal) (M05A) - Additional Information

Gene ID 56997

Other Names

Chaperone activity of bc1 complex-like, mitochondrial, Chaperone-ABC1-like, 2711-, aarF domain-containing protein kinase 3, ADCK3, CABC1

Target/Specificity

CABC1 (AAH05171, 1 a.a. \sim 100 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

CABC1 Antibody (monoclonal) (M05A) is for research use only and not for use in diagnostic or therapeutic procedures.

CABC1 Antibody (monoclonal) (M05A) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot




- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

CABC1 Antibody (monoclonal) (M05A) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa).



CABC1 monoclonal antibody (M05A), clone 8F7 Western Blot analysis of CABC1 expression in Hela S3 NE ((Cat # AT1362a)

CABC1 Antibody (monoclonal) (M05A) - Background

This gene encodes a mitochondrial protein similar to yeast ABC1, which functions in an electron-transferring membrane protein complex in the respiratory chain. It is not related to the family of ABC transporter proteins. Expression of this gene is induced by the tumor suppressor p53 and in response to DNA damage, and inhibiting its expression partially suppresses p53-induced apoptosis. Alternatively spliced transcript variants have been found; however, their full-length nature has not been determined.

CABC1 Antibody (monoclonal) (M05A) - References

ADCK3, an ancestral kinase, is mutated in a form of recessive ataxia associated with coenzyme Q10 deficiency. Lagier-Tourenne C, et al. Am J Hum Genet, 2008 Mar. PMID 18319074.CABC1 gene mutations cause ubiquinone deficiency with cerebellar ataxia and seizures. Mollet J, et al. Am J Hum





Tel: 858.875.1900 Fax: 858.875.1999

Genet, 2008 Mar. PMID 18319072. The DNA sequence and biological annotation of human chromosome 1. Gregory SG, et al. Nature, 2006 May 18. PMID 16710414. Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes. Kimura K, et al. Genome Res, 2006 Jan. PMID 16344560. Towards a proteome-scale map of the human protein-protein interaction network. Rual JF, et al. Nature, 2005 Oct 20. PMID 16189514.