

**CDK4 Antibody (monoclonal) (M04)****Mouse monoclonal antibody raised against a partial recombinant CDK4.****Catalog # AT1475a****Specification**

---

**CDK4 Antibody (monoclonal) (M04) - Product Information**

Application	WB, IF, E
Primary Accession	<a href="#">P11802</a>
Other Accession	<a href="#">BC003644</a>
Reactivity	Human, Mouse
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	33730

**CDK4 Antibody (monoclonal) (M04) - Additional Information****Gene ID** 1019**Other Names**

Cyclin-dependent kinase 4, Cell division protein kinase 4, PSK-J3, CDK4

**Target/Specificity**

CDK4 (AAH03644, 211 a.a. ~ 303 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

IF~~1:50~200

E~~N/A

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

CDK4 Antibody (monoclonal) (M04) is for research use only and not for use in diagnostic or therapeutic procedures.


**CDK4 Antibody (monoclonal) (M04) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### CDK4 Antibody (monoclonal) (M04) - Images




Immunofluorescence of monoclonal antibody to CDK4 on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (35.97 KDa) .




CDK4 monoclonal antibody (M04), clone 2G7 Western Blot analysis of CDK4 expression in HeLa (Cat # AT1475a )



CDK4 monoclonal antibody (M04), clone 2G7. Western Blot analysis of CDK4 expression in Raw 264.7 (Cat # AT1475a )



CDK4 monoclonal antibody (M04), clone 2G7. Western Blot analysis of CDK4 expression in NIH/3T3 (Cat # AT1475a )



Detection limit for recombinant GST tagged CDK4 is approximately 0.1ng/ml as a capture antibody.

#### CDK4 Antibody (monoclonal) (M04) - Background

The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is highly similar to the gene products of *S. cerevisiae* cdc28 and *S. pombe* cdc2. It is a catalytic subunit of the protein kinase complex that is important for cell cycle G1 phase progression. The activity of this kinase is restricted to the G1-S phase, which is controlled by the regulatory subunits D-type cyclins and CDK inhibitor p16(INK4a). This kinase was shown to be responsible for the

phosphorylation of retinoblastoma gene product (Rb). Mutations in this gene as well as in its related proteins including D-type cyclins, p16(INK4a) and Rb were all found to be associated with tumorigenesis of a variety of cancers. Multiple polyadenylation sites of this gene have been reported.

#### **CDK4 Antibody (monoclonal) (M04) - References**

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. The Ras oncogene signals centrosome amplification in mammary epithelial cells through cyclin D1/Cdk4 and Nek2. Zeng X, et al. Oncogene, 2010 Sep 9. PMID 20581865. Pattern of retinoblastoma pathway inactivation dictates response to CDK4/6 inhibition in GBM. Wiedemeyer WR, et al. Proc Natl Acad Sci U S A, 2010 Jun 22. PMID 20534551. A Large-scale genetic association study of esophageal adenocarcinoma risk. Liu CY, et al. Carcinogenesis, 2010 Jul. PMID 20453000. Pharmacologic inhibition of cyclin-dependent kinases 4 and 6 arrests the growth of glioblastoma multiforme intracranial xenografts. Michaud K, et al. Cancer Res, 2010 Apr 15. PMID 20354191.