

## CX3CR1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant CX3CR1. Catalog # AT1687a

#### Specification

## CX3CR1 Antibody (monoclonal) (M01) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW WB, E P49238 BC028078 Human Mouse Monoclonal IgG2a Kappa 40396

## CX3CR1 Antibody (monoclonal) (M01) - Additional Information

Gene ID 1524

**Other Names** CX3C chemokine receptor 1, C-X3-C CKR-1, CX3CR1, Beta chemokine receptor-like 1, CMK-BRL-1, CMK-BRL1, Fractalkine receptor, G-protein coupled receptor 13, V28, CX3CR1, CMKBRL1, GPR13

**Target/Specificity** CX3CR1 (AAH28078, 1 a.a. ~ 36 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions** CX3CR1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

#### CX3CR1 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides




- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

CX3CR1 Antibody (monoclonal) (M01) - Images

175 -	
83 -	
62 -	
47.5=	
32.5=	•
25 -	

Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (29.7 KDa) .



Detection limit for recombinant GST tagged CX3CR1 is approximately 10ng/ml as a capture antibody.

# CX3CR1 Antibody (monoclonal) (M01) - Background

Fractalkine is a transmembrane protein and chemokine involved in the adhesion and migration of leukocytes. The protein encoded by this gene is a receptor for fractalkine. The encoded protein also is a coreceptor for HIV-1, and some variations in this gene lead to increased susceptibility to HIV-1 infection and rapid progression to AIDS. Four transcript variants encoding two different isoforms have been found for this gene.

# CX3CR1 Antibody (monoclonal) (M01) - References

1.The involvement of the fractalkine receptor in the transmigration of neuroblastoma cells through bone-marrow endothelial cells.Nevo I, Sagi-Assif O, Meshel T, Ben-Baruch A, Johrer K, Greil R, Trejo LE, Kharenko O, Feinmesser M, Yron I, Witz IP.Cancer Lett. 2009 Jan 8;273(1):127-39. Epub 2008 Sep 7.