

DNAJB2 Antibody (monoclonal) (M02)**Mouse monoclonal antibody raised against a partial recombinant DNAJB2.****Catalog # AT1789a****Specification**

DNAJB2 Antibody (monoclonal) (M02) - Product Information

Application	WB, E
Primary Accession	P25686
Other Accession	NM_006736
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	35580

DNAJB2 Antibody (monoclonal) (M02) - Additional Information**Gene ID** 3300**Other Names**

DnaJ homolog subfamily B member 2, DnaJ protein homolog 1, Heat shock 40 kDa protein 3, Heat shock protein J1, HSJ-1, DNAJB2, HSJ1, HSPF3

Target/Specificity

DNAJB2 (NP_006727, 216 a.a. ~ 324 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

DNAJB2 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.


DNAJB2 Antibody (monoclonal) (M02) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


DNAJB2 Antibody (monoclonal) (M02) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.73 kDa) .



DNAJB2 monoclonal antibody (M02), clone 1B7. Western Blot analysis of DNAJB2 expression in A-431 (Cat # AT1789a)



Detection limit for recombinant GST tagged DNAJB2 is approximately 0.3ng/ml as a capture antibody.

DNAJB2 Antibody (monoclonal) (M02) - References

Oxidative stress, telomere length and biomarkers of physical aging in a cohort aged 79 years from the 1932 Scottish Mental Survey. Starr JM, et al. Mech Ageing Dev, 2008 Dec. PMID 18977241. Gene expression profiling in human skeletal muscle during recovery from eccentric exercise. Mahoney DJ, et al. Am J Physiol Regul Integr Comp Physiol, 2008 Jun. PMID 18321953. Mutation in WNT10A is associated with an autosomal recessive ectodermal dysplasia: the odonto-onycho-dermal dysplasia. Adaimy L, et al. Am J Hum Genet, 2007 Oct. PMID 17847007. A genetic association analysis of cognitive ability and cognitive ageing using 325 markers for 109 genes associated with oxidative stress or cognition. Harris SE, et al. BMC Genet, 2007 Jul 2. PMID 17601350. The UBL domain of PLIC-1 regulates aggresome formation. Heir R, et al. EMBO Rep, 2006 Dec. PMID 17082820.