

F9 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant F9.****Catalog # AT1981a****Specification****F9 Antibody (monoclonal) (M01) - Product Information**

Application	WB, IP, E
Primary Accession	P00740
Other Accession	NM_000133
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	51778

F9 Antibody (monoclonal) (M01) - Additional Information**Gene ID 2158****Other Names**

Coagulation factor IX, Christmas factor, Plasma thromboplastin component, PTC, Coagulation factor IXa light chain, Coagulation factor IXa heavy chain, F9

Target/Specificity

F9 (NP_000124, 96 a.a. ~ 190 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

IP~~N/A

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

F9 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


F9 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


F9 Antibody (monoclonal) (M01) - Images



Immunoprecipitation of F9 transfected lysate using anti-F9 monoclonal antibody and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with F9 MaxPab rabbit polyclonal antibody.




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.19 KDa) .




Western Blot analysis of F9 expression in transfected 293T cell line by F9 monoclonal antibody (M01), clone 2C9.

Lane 1: F9 transfected lysate(51.8 KDa).

Lane 2: Non-transfected lysate.



Western blot analysis of F9 over-expressed 293 cell line, cotransfected with F9 Validated Chimera RNAi (Cat # AT1981a)



Detection limit for recombinant GST tagged F9 is approximately 0.1ng/ml as a capture antibody.

F9 Antibody (monoclonal) (M01) - Background

This gene encodes vitamin K-dependent coagulation factor IX that circulates in the blood as an

inactive zymogen. This factor is converted to an active form by factor Xla, which excises the activation peptide and thus generates a heavy chain and a light chain held together by one or more disulfide bonds. The role of this activated factor IX in the blood coagulation cascade is to activate factor X to its active form through interactions with Ca+2 ions, membrane phospholipids, and factor VIII. Alterations of this gene, including point mutations, insertions and deletions, cause factor IX deficiency, which is a recessive X-linked disorder, also called hemophilia B or Christmas disease.

F9 Antibody (monoclonal) (M01) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolidinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. *Diabetes Care*, 2010 Jul 13. PMID 20628086. Genetic risk factors for hepatopulmonary syndrome in patients with advanced liver disease. Roberts KE, et al. *Gastroenterology*, 2010 Jul. PMID 20346360. Gene variants associated with venous thrombosis: confirmation in the MEGA study. Arellano AR, et al. *J Thromb Haemost*, 2010 May. PMID 20128871. Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. *Am J Hum Genet*, 2009 Nov. PMID 19913121. Contribution of magnesium in binding of factor IXa to the phospholipid surface: implications for vitamin K-dependent coagulation proteins. Messer AS, et al. *J Thromb Haemost*, 2009 Dec. PMID 19817987.