

## FABP1 Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a full length recombinant FABP1. Catalog # AT1984a

#### Specification

## FABP1 Antibody (monoclonal) (M02) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW WB, E <u>P07148</u> <u>BC032801</u> Human mouse Monoclonal IgG2b Kappa 14208

## FABP1 Antibody (monoclonal) (M02) - Additional Information

Gene ID 2168

**Other Names** Fatty acid-binding protein, liver, Fatty acid-binding protein 1, Liver-type fatty acid-binding protein, L-FABP, FABP1, FABPL

**Target/Specificity** FABP1 (AAH32801, 1 a.a. ~ 127 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution** WB~~1:500~1000 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions** FABP1 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.

#### FABP1 Antibody (monoclonal) (M02) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides



- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- <u>Flow Cytomety</u>
- <u>Cell Culture</u>

FABP1 Antibody (monoclonal) (M02) - Images



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (39.71 KDa).



Western Blot analysis of FABP1 expression in transfected 293T cell line by FABP1 monoclonal antibody (M02), clone 5F7.

Lane 1: FABP1 transfected lysate(14.2 KDa). Lane 2: Non-transfected lysate.





Western blot analysis of FABP1 over-expressed 293 cell line, cotransfected with FABP1 Validated Chimera RNAi ( (Cat # AT1984a )



Detection limit for recombinant GST tagged FABP1 is approximately 0.1ng/ml as a capture antibody.

# FABP1 Antibody (monoclonal) (M02) - Background

FABP1 encodes the fatty acid binding protein found in liver. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that bind long-chain fatty acids and other hydrophobic ligands. FABP1 and FABP6 (the ileal fatty acid binding protein) are also able to bind bile acids. It is thought that FABPs roles include fatty acid uptake, transport, and metabolism.

# FABP1 Antibody (monoclonal) (M02) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.Millisecond timescale dynamics of human liver fatty acid binding protein: testing of its relevance to the ligand entry process. Long D, et al. Biophys J, 2010 Jun 16. PMID 20550918.Identification of type 2 diabetes-associated combination of SNPs using support vector machine. Ban HJ, et al. BMC Genet, 2010 Apr 23. PMID 20416077.Urinary liver-type fatty acid-binding protein predicts progression to nephropathy in type 1 diabetic patients. Nielsen SE, et al. Diabetes Care, 2010 Jun. PMID 20185732.Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121.