

### FBXO8 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant FBXO8. Catalog # AT2022a

## **Specification**

## FBXO8 Antibody (monoclonal) (M01) - Product Information

**Application** WB, E **Primary Accession** O9NRD0 NM 012180 Other Accession Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG1 Kappa Calculated MW 37068

## FBXO8 Antibody (monoclonal) (M01) - Additional Information

#### **Gene ID 26269**

#### **Other Names**

F-box only protein 8, F-box/SEC7 protein FBS, FBXO8, FBS, FBX8

# **Target/Specificity**

FBXO8 (NP\_036312, 1 a.a.  $\sim$  77 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

## **Dilution**

WB~~1:500~1000

#### **Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2.

#### **Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

### **Precautions**

FBXO8 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

## FBXO8 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry




- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## FBXO8 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (34.21 KDa).



Detection limit for recombinant GST tagged FBXO8 is approximately 0.1ng/ml as a capture antibody.

## FBXO8 Antibody (monoclonal) (M01) - Background

This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of the ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbxs class. It contains a C-terminal amino acid sequence that bears a significant similarity with a portion of yeast Sec7p, a critical regulator of vesicular protein transport. This human protein may interact with ADP-ribosylation factor(s)(ARFs) and exhibit ARF-GEF (guanine nucleotide exchange factor) activity.

## FBXO8 Antibody (monoclonal) (M01) - References

Screening for replication of genome-wide SNP associations in sporadic ALS. Cronin S, et al. Eur J Hum Genet, 2009 Feb. PMID 18987618. Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome







Res, 2004 Oct. PMID 15489334. Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. Nat Genet, 2004 Jan. PMID 14702039. The secreted protein discovery initiative (SPDI), a large-scale effort to identify novel human secreted and transmembrane proteins: a bioinformatics assessment. Clark HF, et al. Genome Res, 2003 Oct. PMID 12975309.