

GRIA1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant GRIA1. Catalog # AT2261a

Specification

GRIA1 Antibody (monoclonal) (M01) - Product Information

Application WB, E **Primary Accession** P42261 Other Accession NM 000827 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG2a Kappa Calculated MW 101506

GRIA1 Antibody (monoclonal) (M01) - Additional Information

Gene ID 2890

Other Names

Glutamate receptor 1, GluR-1, AMPA-selective glutamate receptor 1, GluR-A, GluR-K1, Glutamate receptor ionotropic, AMPA 1, GluA1, GRIA1, GLUH1, GLUR1

Target/Specificity

GRIA1 (NP_000818, 201 a.a. \sim 300 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000 E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

GRIA1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

GRIA1 Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides




- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

GRIA1 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa).



Detection limit for recombinant GST tagged GRIA1 is approximately 0.3ng/ml as a capture antibody.

GRIA1 Antibody (monoclonal) (M01) - Background

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. These receptors are heteromeric protein complexes with multiple subunits, each possessing transmembrane regions, and all arranged to form a ligand-gated ion channel. The classification of glutamate receptors is based on their activation by different pharmacologic agonists. This gene belongs to a family of alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate (AMPA) receptors. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

GRIA1 Antibody (monoclonal) (M01) - References

Genetic variations in GRIA1 on chromosome 5q33 related to asparaginase hypersensitivity. Chen SH, et al. Clin Pharmacol Ther, 2010 Aug. PMID 20592726.Common variants in the regulative regions of GRIA1 and GRIA3 receptor genes are associated with migraine susceptibility. Formicola D, et al. BMC Med Genet, 2010 Jun 25. PMID 20579352.Association study of 182 candidate genes in anorexia nervosa. Pinheiro AP, et al. Am J Med Genet B Neuropsychiatr Genet, 2010 Jul. PMID







20468064. Comprehensive copy number variant (CNV) analysis of neuronal pathways genes in psychiatric disorders identifies rare variants within patients. Saus E, et al. J Psychiatr Res, 2010 Apr 14. PMID 20398908. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.