

**HARS Antibody (monoclonal) (M01)****Mouse monoclonal antibody raised against a partial recombinant HARS.****Catalog # AT2317a****Specification**

---

**HARS Antibody (monoclonal) (M01) - Product Information**

Application	WB, IF, E
Primary Accession	<a href="#">P12081</a>
Other Accession	<a href="#">NM_002109</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	57411

**HARS Antibody (monoclonal) (M01) - Additional Information****Gene ID** 3035**Other Names**

Histidine--tRNA ligase, cytoplasmic, Histidyl-tRNA synthetase, HisRS, HARS, HRS

**Target/Specificity**

HARS (NP\_002100, 1 a.a. ~ 96 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

IF~~1:50~200

E~~N/A

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

HARS Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


**HARS Antibody (monoclonal) (M01) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### HARS Antibody (monoclonal) (M01) - Images




Immunofluorescence of monoclonal antibody to HARS on HeLa cell. [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.3 KDa) .



HARS monoclonal antibody (M01), clone 1C8 Western Blot analysis of HARS expression in HeLa (Cat # AT2317a )



Detection limit for recombinant GST tagged HARS is approximately 0.03ng/ml as a capture antibody.

### **HARS Antibody (monoclonal) (M01) - Background**

Aminoacyl-tRNA synthetases are a class of enzymes that charge tRNAs with their cognate amino acids. The protein encoded by this gene is a cytoplasmic enzyme which belongs to the class II family of aminoacyl-tRNA synthetases. The enzyme is responsible for the synthesis of histidyl-transfer RNA, which is essential for the incorporation of histidine into proteins. The gene is located in a head-to-head orientation with HARS1 on chromosome five, where the homologous genes share a bidirectional promoter. The gene product is a frequent target of autoantibodies in the human autoimmune disease polymyositis/dermatomyositis.

### **HARS Antibody (monoclonal) (M01) - References**

Clinical significance of anti-histidyl-tRNA synthetase (Jo1) autoantibodies. Gomard-Mennesson E, et al. Ann N Y Acad Sci, 2007 Aug. PMID 17785330. Novel conformation of histidyl-transfer RNA synthetase in the lung: the target tissue in Jo-1 autoantibody-associated myositis. Levine SM, et al. Arthritis Rheum, 2007 Aug. PMID 17665459. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334. Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. Nat Genet, 2004 Jan. PMID 14702039. TSG101 interaction with HRS mediates endosomal trafficking and receptor down-regulation. Lu Q, et al. Proc Natl Acad Sci U S A, 2003 Jun 24. PMID 12802020.