

HFE Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant HFE. Catalog # AT2361a

Specification

HFE Antibody (monoclonal) (M01) - Product Information

Application IF, WB, E **Primary Accession** 030201 Other Accession NM 000410 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG1 Kappa Calculated MW 40108

HFE Antibody (monoclonal) (M01) - Additional Information

Gene ID 3077

Other Names

Hereditary hemochromatosis protein, HLA-H, HFE, HLAH

Target/Specificity

HFE (NP_000401, 115 a.a. \sim 205 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

HFE Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

HFE Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry




- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture


HFE Antibody (monoclonal) (M01) - Images



Immunofluorescence of monoclonal antibody to HFE on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (35.75 KDa).



HFE monoclonal antibody (M01), clone 1G12. Western Blot analysis of HFE expression in A-431.






Western Blot analysis of HFE expression in transfected 293T cell line by HFE monoclonal antibody (M01), clone 1G12.

Lane 1: HFE transfected lysate(40.1 KDa).

Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged HFE is approximately 0.03ng/ml as a capture antibody.

HFE Antibody (monoclonal) (M01) - Background

The protein encoded by this gene is a membrane protein that is similar to MHC class I-type proteins and associates with beta2-microglobulin (beta2M). It is thought that this protein functions to regulate iron absorption by regulating the interaction of the transferrin receptor with transferrin. The iron storage disorder, hereditary haemochromatosis, is a recessive genetic disorder that results from defects in this gene. At least nine alternatively spliced variants have been described for this gene. Additional variants have been found but their full-length nature has not been determined.

HFE Antibody (monoclonal) (M01) - References

1.Hemochromatosis Enhances Tumor Progression via Upregulation of Intracellular Iron in Head and Neck Cancer.Lenarduzzi M, Hui AB, Yue S, Ito E, Shi W, Williams J, Bruce J, Sakemura-Nakatsugawa N, Xu W, Schimmer A, Liu FFPLoS One. 2013 Aug 26;8(8):e74075. doi: 10.1371/journal.pone.0074075.