10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HHEX Antibody (monoclonal) (M05)

Mouse monoclonal antibody raised against a full length recombinant HHEX.
Catalog # AT2365a

Specification

HHEX Antibody (monoclonal) (M05) - Product Information

Application	**WB, E**
Primary Accession	Q03014
Other Accession	BC014336
Reactivity	**Human**
Host	**mouse**
Clonality	**Monoclonal**
Isotype	**IgG2a Lambda**
Calculated MW	**30022**

HHEX Antibody (monoclonal) (M05) - Additional Information

Gene ID 3087

Other Names
Hematopoietically-expressed homeobox protein HHEX, Homeobox protein HEX, Homeobox protein PRH, HHEX, HEX, PRH, PRHX

Target/Specificity
HHEX (AAH14336, 1 a.a. ~ 270 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution
WB~~1:500~1000
E~~N/A

Format
Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage
Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions
HHEX Antibody (monoclonal) (M05) is for research use only and not for use in diagnostic or therapeutic procedures.

HHEX Antibody (monoclonal) (M05) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

HHEX Antibody (monoclonal) (M05) - Images



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (55.44 KDa) .



Detection limit for recombinant GST tagged HHEX is approximately 0.1ng/ml as a capture antibody.

HHEX Antibody (monoclonal) (M05) - Background

This gene encodes a member of the homeobox family of transcription factors, many of which are involved in developmental processes. Expression in specific hematopoietic lineages suggests that this protein may play a role in hematopoietic differentiation.

HHEX Antibody (monoclonal) (M05) - References

Glycemia determines the effect of type 2 diabetes risk genes on insulin secretion. Heni M, et al. Diabetes, 2010 Aug 29. PMID 20802253.Association between variants in IDE-KIF11-HHEX and plasma amyloid beta levels. Reitz C, et al. Neurobiol Aging, 2010 Aug 17. PMID 20724036.No association between FTO or HHEX and endometrial cancer risk. Gaudet MM, et al. Cancer Epidemiol Biomarkers Prev, 2010 Aug. PMID 20647405.Coronary artery calcification and its relationship to validated genetic variants for diabetes mellitus assessed in the Heinz Nixdorf recall cohort. Pechlivanis S, et al. Arterioscler Thromb Vasc Biol, 2010 Sep. PMID 20616309.A validation study of type 2 diabetes-related variants of the TCF7L2, HHEX, KCNJ11, and ADIPOQ genes in one endogamous ethnic group of north India. Gupta V, et al. Ann Hum Genet, 2010 Jul. PMID 20597906.