

IFI16 Antibody (monoclonal) (M03)**Mouse monoclonal antibody raised against a partial recombinant IFI16.****Catalog # AT2484a****Specification****IFI16 Antibody (monoclonal) (M03) - Product Information**

Application	IF, WB, IHC, E
Primary Accession	Q16666
Other Accession	BC017059
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b Kappa
Calculated MW	88256

IFI16 Antibody (monoclonal) (M03) - Additional Information**Gene ID** 3428**Other Names**

Gamma-interferon-inducible protein 16, Ifi-16, Interferon-inducible myeloid differentiation transcriptional activator, IFI16, IFNGIP1

Target/Specificity

IFI16 (AAH17059, 630 a.a. ~ 729 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions


IFI16 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.

IFI16 Antibody (monoclonal) (M03) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)


- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IFI16 Antibody (monoclonal) (M03) - Images

Immunofluorescence of monoclonal antibody to IFI16 on HeLa cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .



IFI16 monoclonal antibody (M03), clone 2E3 Western Blot analysis of IFI16 expression in Hela S3


NE ((Cat # AT2484a)




Western Blot analysis of IFI16 expression in transfected 293T cell line by IFI16 monoclonal antibody (M03), clone 2E3.

Lane 1: IFI16 transfected lysate(82 KDa).

Lane 2: Non-transfected lysate.



Immunoperoxidase of monoclonal antibody to IFI16 on formalin-fixed paraffin-embedded human tonsil. [antibody concentration 3 ug/ml]



Detection limit for recombinant GST tagged IFI16 is approximately 0.3ng/ml as a capture antibody.

IFI16 Antibody (monoclonal) (M03) - Background

This gene encodes a member of the HIN-200 (hematopoietic interferon-inducible nuclear antigens with 200 amino acid repeats) family of cytokines. The encoded protein contains domains involved in

DNA binding, transcriptional regulation, and protein-protein interactions. The protein localizes to the nucleoplasm and nucleoli, and interacts with p53 and retinoblastoma-1. It modulates p53 function, and inhibits cell growth in the Ras/Raf signaling pathway.

IFI16 Antibody (monoclonal) (M03) - References

Evaluation of candidate stromal epithelial cross-talk genes identifies association between risk of serous ovarian cancer and TERT, a cancer susceptibility hot-spot. Johnatty SE, et al. PLoS Genet, 2010 Jul 8. PMID 20628624. Association of IFI200 gene polymorphisms with susceptibility to systemic lupus erythematosus. Kimkong I, et al. J Rheumatol, 2010 Jul. PMID 20595294. Human cytomegalovirus pUL83 stimulates activity of the viral immediate-early promoter through its interaction with the cellular IFI16 protein. Cristea IM, et al. J Virol, 2010 Aug. PMID 20504932. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. Delta-like 1 contributes to cell growth by increasing the interferon-inducible protein 16 expression in hepatocellular carcinoma. Yu F, et al. Liver Int, 2010 May. PMID 20214740.