

IQGAP2 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant IQGAP2. Catalog # AT2536a

Specification

IQGAP2 Antibody (monoclonal) (M01) - Product Information

Application WB, E **Primary Accession** 013576 Other Accession NM 006633 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG2b Kappa Calculated MW 180578

IQGAP2 Antibody (monoclonal) (M01) - Additional Information

Gene ID 10788

Other Names

Ras GTPase-activating-like protein IQGAP2, IQGAP2

Target/Specificity

IQGAP2 (NP_006624, 343 a.a. \sim 449 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

IQGAP2 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

IQGAP2 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot




- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


IQGAP2 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.51 KDa).



IQGAP2 monoclonal antibody (M01), clone 1B10 Western Blot analysis of IQGAP2 expression in HeLa ((Cat # AT2536a)



Detection limit for recombinant GST tagged IQGAP2 is approximately 0.03ng/ml as a capture antibody.

IQGAP2 Antibody (monoclonal) (M01) - Background





Tel: 858.875.1900 Fax: 858.875.1999

This gene encodes a member of the IQGAP family. The protein contains three IQ domains, one calponin homology domain, one Ras-GAP domain and one WW domain. It interacts with components of the cytoskeleton, with cell adhesion molecules, and with several signaling molecules to regulate cell morphology and motility.

IQGAP2 Antibody (monoclonal) (M01) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.A genome-wide association study for age-related hearing impairment in the Saami. Van Laer L, et al. Eur J Hum Genet, 2010 Jun. PMID 20068591. Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121.IQGAP2 inactivation through aberrant promoter methylation and promotion of invasion in gastric cancer cells. Jin SH, et al. Int J Cancer, 2008 Mar 1. PMID 17957782.