

KCNJ15 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant KCNJ15. Catalog # AT2599a

Specification

KCNJ15 Antibody (monoclonal) (M01) - Product Information

WB, E Application **Primary Accession** 099712 Other Accession NM 002243 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG2a Kappa Calculated MW 42577

KCNJ15 Antibody (monoclonal) (M01) - Additional Information

Gene ID 3772

Other Names

ATP-sensitive inward rectifier potassium channel 15, Inward rectifier K(+) channel Kir13, Inward rectifier K(+) channel Kir42, Potassium channel, inwardly rectifying subfamily J member 15, KCNJ14

Target/Specificity

KCNJ15 (NP_002234, 290 a.a. \sim 355 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

KCNJ15 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

KCNJ15 Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides




- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

KCNJ15 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (33 KDa).




Western Blot analysis of KCNJ15 expression in transfected 293T cell line by KCNJ15 monoclonal antibody (M01), clone 1B2.


Lane 1: KCNJ15 transfected lysate(42.6 KDa).

Lane 2: Non-transfected lysate.





Western blot analysis of KCNJ15 over-expressed 293 cell line, cotransfected with KCNJ15 Validated Chimera RNAi ((Cat # AT2599a)



Detection limit for recombinant GST tagged KCNJ15 is 3 ng/ml as a capture antibody.

KCNJ15 Antibody (monoclonal) (M01) - Background

Potassium channels are present in most mammalian cells, where they participate in a wide range of physiologic responses. The protein encoded by this gene is an integral membrane protein and inward-rectifier type potassium channel. The encoded protein has a greater tendency to allow potassium to flow into a cell rather than out of a cell. Three transcript variants encoding the same protein have been found for this gene.

KCNJ15 Antibody (monoclonal) (M01) - References

Identification of KCNJ15 as a susceptibility gene in Asian patients with type 2 diabetes mellitus. Okamoto K, et al. Am J Hum Genet, 2010 Jan. PMID 20085713.MUPP1 complexes renal K+ channels to alter cell surface expression and whole cell currents. Sindic A, et al. Am J Physiol Renal Physiol, 2009 Jul. PMID 19420109.Rare independent mutations in renal salt handling genes contribute to blood pressure variation. Ji W, et al. Nat Genet, 2008 May. PMID 18391953.Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348.Interaction of the Ca2+-sensing receptor with the inwardly rectifying potassium channels Kir4.1 and Kir4.2 results in inhibition of channel function. Huang C, et al. Am J Physiol Renal Physiol, 2007 Mar. PMID 17122384.