

LDB3 Antibody (monoclonal) (M06)

Mouse monoclonal antibody raised against a full-length recombinant LDB3. Catalog # AT2686a

Specification

LDB3 Antibody (monoclonal) (M06) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW WB, IF, E <u>075112</u> <u>BC010929</u> Human mouse Monoclonal IgG2b Kappa 77135

LDB3 Antibody (monoclonal) (M06) - Additional Information

Gene ID 11155

```
Other Names
LIM domain-binding protein 3, Protein cypher, Z-band alternatively spliced PDZ-motif protein, LDB3
(<a href="http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=15710"
target="_blank">HGNC:15710</a>)
```

Target/Specificity LDB3 (AAH10929, 1 a.a. ~ 283 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution WB~~1:500~1000 IF~~1:50~200 E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions LDB3 Antibody (monoclonal) (M06) is for research use only and not for use in diagnostic or therapeutic procedures.


LDB3 Antibody (monoclonal) (M06) - Protocols

Provided below are standard protocols that you may find useful for product applications.




- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>


LDB3 Antibody (monoclonal) (M06) - Images




Immunofluorescence of monoclonal antibody to LDB3 on A-431 cell . [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein.Western Blot detection against Immunogen (56.87 KDa).




LDB3 monoclonal antibody (M06), clone 3C8 Western Blot analysis of LDB3 expression in A-431 ((Cat # AT2686a)



Western Blot analysis of LDB3 expression in transfected 293T cell line by LDB3 monoclonal antibody (M06), clone 3C8.


Lane 1: LDB3 transfected lysate(31 KDa). Lane 2: Non-transfected lysate.



Western blot analysis of LDB3 over-expressed 293 cell line, cotransfected with LDB3 Validated



Chimera RNAi ((Cat # AT2686a)



Detection limit for recombinant GST tagged LDB3 is approximately 0.03ng/ml as a capture antibody.

LDB3 Antibody (monoclonal) (M06) - Background

This gene encodes a PDZ domain-containing protein. PDZ motifs are modular protein-protein interaction domains consisting of 80-120 amino acid residues. PDZ domain-containing proteins interact with each other in cytoskeletal assembly or with other proteins involved in targeting and clustering of membrane proteins. The protein encoded by this gene interacts with alpha-actinin-2 through its N-terminal PDZ domain and with protein kinase C via its C-terminal LIM domains. The LIM domain is a cysteine-rich motif defined by 50-60 amino acids containing two zinc-binding modules. This protein also interacts with all three members of the myozenin family. Mutations in this gene have been associated with myofibrillar myopathy and dilated cardiomyopathy. Alternatively spliced transcript variants encoding different isoforms have been identified; all isoforms have N-terminal PDZ domains while only longer isoforms (1, 2 and 5) have C-terminal LIM domains.

LDB3 Antibody (monoclonal) (M06) - References

1.Z-disc-associated, Alternatively Spliced, PDZ Motif-containing Protein (ZASP) Mutations in the Actin-binding Domain Cause Disruption of Skeletal Muscle Actin Filaments in Myofibrillar Myopathy.Lin X, Ruiz J, Bajraktari I, Ohman R, Banerjee S, Gribble K, Kaufman JD, Wingfield PT, Griggs RC, Fischbeck KH, Mankodi AJ Biol Chem. 2014 May 9;289(19):13615-26. doi: 10.1074/jbc.M114.550418. Epub 2014 Mar 25.2.Impaired binding of ZASP/Cypher with phosphoglucomutase 1 is associated with dilated cardiomyopathy.Arimura T, Inagaki N, Hayashi T, Shichi D, Sato A, Hinohara K, Vatta M, Towbin JA, Chikamori T, Yamashina A, Kimura A.Cardiovasc Res. 2009 Jul 1;83(1):80-8. Epub 2009 Apr 17.