

MAP3K5 Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a partial recombinant MAP3K5. Catalog # AT2777a

Specification

MAP3K5 Antibody (monoclonal) (M03) - Product Information

WB, IF, E Application **Primary Accession** 099683 Other Accession BC054503 Reactivity Human Host mouse Clonality **Monoclonal** Isotype **IgG1 Lambda** Calculated MW 154537

MAP3K5 Antibody (monoclonal) (M03) - Additional Information

Gene ID 4217

Other Names

Mitogen-activated protein kinase kinase 5, Apoptosis signal-regulating kinase 1, ASK-1, MAPK/ERK kinase kinase 5, MEK kinase 5, MEKK 5, MAP3K5, ASK1, MAPKKK5

Target/Specificity

MAP3K5 (AAH54503, 1231 a.a. \sim 1374 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000 IF~~1:50~200 E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

MAP3K5 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.

MAP3K5 Antibody (monoclonal) (M03) - Protocols


Provided below are standard protocols that you may find useful for product applications.

• Western Blot




- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


MAP3K5 Antibody (monoclonal) (M03) - Images



Immunofluorescence of monoclonal antibody to MAP3K5 on A-431 cell. [antibody concentration 10 ug/ml]




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen $(41.58\ \text{KDa})$.






MAP3K5 monoclonal antibody (M03), clone 2D11 Western Blot analysis of MAP3K5 expression in A-431 ((Cat # AT2777a)



Western Blot analysis of MAP3K5 expression in transfected 293T cell line by MAP3K5 monoclonal antibody (M03), clone 2D11.

Lane 1: MAP3K5 transfected lysate(155 KDa).

Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged MAP3K5 is approximately 0.3ng/ml as a capture antibody.

MAP3K5 Antibody (monoclonal) (M03) - Background

Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulated kinase (ERK), MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAPKKK or MEKK). MAPKK kinase/MEKK phosphorylates and activates its downstream protein kinase, MAPK kinase/MEK, which in turn activates MAPK. The kinases of these signaling cascades are highly conserved, and homologs exist in yeast, Drosophila, and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. Northern blot analysis shows that MAPKKK5 transcript is abundantly expressed in human heart and pancreas. The MAPKKK5 protein phosphorylates and activates MKK4 (aliases SERK1, MAPKK4) in vitro, and activates c-Jun N-terminal kinase (JNK)/stress-activated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK.

MAP3K5 Antibody (monoclonal) (M03) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.ASK1-JNK signaling cascade mediates





Tel: 858.875.1900 Fax: 858.875.1999

Ad-ST13-induced apoptosis in colorectal HCT116 cells. Yang M, et al. J Cell Biochem, 2010 Jun 1. PMID 20512919. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. Mechanism of oxidative stress-induced ASK1-catalyzed MKK6 phosphorylation. Sturchler E, et al. Biochemistry, 2010 May 18. PMID 20364819.DJ-1 modulates the p38 mitogen-activated protein kinase pathway through physical interaction with apoptosis signal-regulating kinase 1. Mo JS, et al. J Cell Biochem, 2010 May. PMID 20213747.