

MAZ Antibody (monoclonal) (M05)**Mouse monoclonal antibody raised against a partial recombinant MAZ.****Catalog # AT2809a****Specification**

MAZ Antibody (monoclonal) (M05) - Product Information

Application	WB, E
Primary Accession	P56270
Other Accession	NM_002383
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	48608

MAZ Antibody (monoclonal) (M05) - Additional Information**Gene ID** 4150**Other Names**

Myc-associated zinc finger protein, MAZI, Pur-1, Purine-binding transcription factor, Serum amyloid A-activating factor-1, SAF-1, Transcription factor Zif87, ZF87, Zinc finger protein 801, MAZ, ZNF801

Target/Specificity

MAZ (NP_002374, 332 a.a. ~ 440 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

MAZ Antibody (monoclonal) (M05) is for research use only and not for use in diagnostic or therapeutic procedures.


MAZ Antibody (monoclonal) (M05) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MAZ Antibody (monoclonal) (M05) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.73 KDa) .



Detection limit for recombinant GST tagged MAZ is approximately 0.1ng/ml as a capture antibody.

MAZ Antibody (monoclonal) (M05) - References

Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732. SAF-3, a novel splice variant of the SAF-1/MAZ/Pur-1 family, is expressed during inflammation. Ray A, et al. FEBS J, 2009 Aug. PMID 19583771. Association and mutation analyses of 16p11.2 autism candidate genes. Kumar RA, et al. PLoS One, 2009. PMID 19242545. Gastrin activates paracrine networks leading to induction of PAI-2 via MAZ and ASC-1. Almeida-Vega S, et al. Am J Physiol Gastrointest Liver Physiol, 2009 Feb. PMID 19074642. Transcriptional synergy mediated by SAF-1 and AP-1: critical role of N-terminal polyalanine and two zinc finger domains of SAF-1. Kumar D, et al. J Biol Chem, 2009 Jan 16. PMID 19028685.