

MBD1 Antibody (monoclonal) (M05)**Mouse monoclonal antibody raised against a partial recombinant MBD1.****Catalog # AT2812a****Specification**

MBD1 Antibody (monoclonal) (M05) - Product Information

Application	WB, E
Primary Accession	O9UIS9
Other Accession	NM_015846
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2b Kappa
Calculated MW	66607

MBD1 Antibody (monoclonal) (M05) - Additional Information**Gene ID** 4152**Other Names**

Methyl-CpG-binding domain protein 1, CXXC-type zinc finger protein 3, Methyl-CpG-binding protein MBD1, Protein containing methyl-CpG-binding domain 1, MBD1, CXXC3, PCM1

Target/Specificity

MBD1 (NP_056671, 415 a.a. ~ 508 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

MBD1 Antibody (monoclonal) (M05) is for research use only and not for use in diagnostic or therapeutic procedures.


MBD1 Antibody (monoclonal) (M05) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


MBD1 Antibody (monoclonal) (M05) - Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.08 KDa) .



MBD1 monoclonal antibody (M05), clone 2B10. Western Blot analysis of MBD1 expression in human kidney.



MBD1 monoclonal antibody (M05), clone 2B10. Western Blot analysis of MBD1 expression in Jurkat (Cat # AT2812a)



Detection limit for recombinant GST tagged MBD1 is 0.3 ng/ml as a capture antibody.

MBD1 Antibody (monoclonal) (M05) - Background

DNA methylation is the major modification of eukaryotic genomes and plays an essential role in mammalian development. Human proteins MECP2, MBD1, MBD2, MBD3, and MBD4 comprise a family of nuclear proteins related by the presence in each of a methyl-CpG binding domain (MBD). Each of these proteins, with the exception of MBD3, is capable of binding specifically to methylated DNA. MECP2, MBD1 and MBD2 can also repress transcription from methylated gene promoters. Five transcript variants of the MBD1 are generated by alternative splicing resulting in protein isoforms that contain one MBD domain, two to three cysteine-rich (CXXC) domains, and some differences in the COOH terminus. All five transcript variants repress transcription from methylated promoters; in addition, variants with three CXXC domains also repress unmethylated promoter activity. MBD1 and MBD2 map very close to each other on chromosome 18q21.

MBD1 Antibody (monoclonal) (M05) - References

Common variants at 2q37.3, 8q24.21, 15q21.3 and 16q24.1 influence chronic lymphocytic leukemia risk. Crowther-Swanepoel D, et al. Nat Genet, 2010 Feb. PMID 20062064. Human BAHD1 promotes heterochromatic gene silencing. Bierne H, et al. Proc Natl Acad Sci U S A, 2009 Aug 18. PMID 19666599. Methyl-CpG binding domain 1 gene polymorphisms and lung cancer risk in a Chinese population. Liu H, et al. Biomarkers, 2008 Sep. PMID 18668384. Silencing of MBD1 and MeCP2 in prostate-cancer-derived PC3 cells produces differential gene expression profiles and cellular phenotypes. Yaqinuddin A, et al. Biosci Rep, 2008 Dec. PMID 18666890. Proteomic analysis of differential proteins in pancreatic carcinomas: Effects of MBD1 knock-down by stable RNA interference. Liu C, et al. BMC Cancer, 2008 Apr 29. PMID 18445260.