

MRC1 Antibody (monoclonal) (M02)**Mouse monoclonal antibody raised against a partial recombinant MRC1.****Catalog # AT2899a****Specification**

MRC1 Antibody (monoclonal) (M02) - Product Information

Application	WB, IHC, E
Primary Accession	P22897
Other Accession	NM_002438
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	166012

MRC1 Antibody (monoclonal) (M02) - Additional Information**Gene ID** 4360**Other Names**

Macrophage mannose receptor 1, MMR, C-type lectin domain family 13 member D, C-type lectin domain family 13 member D-like, Macrophage mannose receptor 1-like protein 1, CD206, MRC1, CLEC13D, CLEC13DL, MRC1L1

Target/Specificity

MRC1 (NP_002429, 22 a.a. ~ 130 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

IHC~~1:100~500

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions


MRC1 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.

MRC1 Antibody (monoclonal) (M02) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


MRC1 Antibody (monoclonal) (M02) - Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.73 KDa) .



MRC1 monoclonal antibody (M02), clone 5C11. Western Blot analysis of MRC1 expression in human pancreas.



Immunoperoxidase of monoclonal antibody to MRC1 on formalin-fixed paraffin-embedded human placenta. [antibody concentration 1.5 ug/ml]



Detection limit for recombinant GST tagged MRC1 is approximately 0.03ng/ml as a capture antibody.

MRC1 Antibody (monoclonal) (M02) - Background

The recognition of complex carbohydrate structures on glycoproteins is an important part of several biological processes, including cell-cell recognition, serum glycoprotein turnover, and neutralization of pathogens. The protein encoded by this gene is a type I membrane receptor that mediates the endocytosis of glycoproteins by macrophages. The protein has been shown to bind high-mannose structures on the surface of potentially pathogenic viruses, bacteria, and fungi so that they can be neutralized by phagocytic engulfment. This gene is in close proximity to MRC1L1. The gene loci including this gene, MRC1L1, as well as LOC340843 and LOC340893, consist of two nearly identical, tandemly linked genomic regions, which are thought to be a part of a duplicated region.

MRC1 Antibody (monoclonal) (M02) - References

1. Increased malignancy of oral squamous cell carcinomas (oscc) is associated with macrophage polarization in regional lymph nodes - an immunohistochemical study. Wehrhan F, Buttner-Herold M, Hyckel P, Moebius P, Preidl R, Distel L, Ries J, Amann K, Schmitt C, Neukam FW, Weber M BMC Cancer. 2014 Jul 21;14:522. doi: 10.1186/1471-2407-14-522.
2. Alternative activation of laser-captured murine hemophagocytes. Canna SW, Costa-Reis P, Bernal WE, Chu N, Sullivan KE, Paessler ME, Behrens E M Arthritis Rheumatol. 2014 Jan 27. doi: 10.1002/art.38379.
3. Small oral squamous cell carcinomas with nodal lymphogenic metastasis show increased infiltration of M2 polarized macrophages - an immunohistochemical analysis. Weber M, Buttner-Herold M, Hyckel P, Moebius P, Distel L, Ries J, Amann K, Neukam FW, Wehrhan F. Journal of Cranio-Maxillofacial Surgery(2014), doi: 10.1016/j.jcms.2014.01.035.
4. Xanthogranulomatous cholecystitis: A clinicopathological study of its association with gallbladder carcinoma. Zhuang PY, Zhu MJ, Wang JD, Zhou XP, Quan ZW, Shen JJ Dig Dis. 2012 Sep 21. doi: 10.1111/j.1751-2980.2012.00645.x. [Epub ahead of print]
5. Coronary Atherosclerosis Is Associated With Macrophage Polarization in Epicardial

Adipose Tissue. Hirata Y, Tabata M, Kurobe H, Motoki T, Akaike M, Nishio C, Higashida M, Mikasa H, Nakaya Y, Takanashi S, Igarashi T, Kitagawa T, Sata M. J Am Coll Cardiol. 2011 Jul 12;58(3):248-55.

MRC1 Antibody (monoclonal) (M02) - Citations

- [Characterizing the immune microenvironment in high-risk ductal carcinoma in situ of the breast.](#)