

### MSR1 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant MSR1. Catalog # AT2919a

### **Specification**

### MSR1 Antibody (monoclonal) (M01) - Product Information

Application WB, E **Primary Accession** P21757 Other Accession NM 138715 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG3 Kappa Calculated MW 49762

### MSR1 Antibody (monoclonal) (M01) - Additional Information

#### **Gene ID 4481**

#### **Other Names**

Macrophage scavenger receptor types I and II, Macrophage acetylated LDL receptor I and II, Scavenger receptor class A member 1, CD204, MSR1, SCARA1

### Target/Specificity

MSR1 (NP\_619729, 121 a.a.  $\sim$  220 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

### **Dilution**

WB~~1:500~1000 E~~N/A

# **Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2.

#### Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

# **Precautions**

MSR1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

# MSR1 Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides




- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## MSR1 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa).




Detection limit for recombinant GST tagged MSR1 is approximately 0.3ng/ml as a capture antibody.

## MSR1 Antibody (monoclonal) (M01) - Background

This gene encodes the class A macrophage scavenger receptors, which include three different types (1, 2, 3) generated by alternative splicing of this gene. These receptors or isoforms are macrophage-specific trimeric integral membrane glycoproteins and have been implicated in many macrophage-associated physiological and pathological processes including atherosclerosis, Alzheimer's disease, and host defense. The isoforms type 1 and type 2 are functional receptors and are able to mediate the endocytosis of modified low density lipoproteins (LDLs). The isoform type 3 does not internalize modified LDL (acetyl-LDL) despite having the domain shown to mediate this function in the types 1 and 2 isoforms. It has an altered intracellular processing and is trapped within the endoplasmic reticulum, making it unable to perform endocytosis. The isoform type 3 can inhibit the function of isoforms type 1 and type 2 when co-expressed, indicating a dominant negative effect and suggesting a mechanism for regulation of scavenger receptor activity in macrophages.

## MSR1 Antibody (monoclonal) (M01) - References





Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Common polymorphisms in ITGA2, PON1 and THBS2 are associated with coronary atherosclerosis in a candidate gene association study of the Chinese Han population. Wang Y, et al. J Hum Genet, 2010 Aug. PMID 20485444.CD36 and macrophage scavenger receptor a modulate foam cell formation via inhibition of lipid-laden platelet phagocytosis. Seizer P, et al. Semin Thromb Hemost, 2010 Mar. PMID 20414830. Genetic variation in APOJ, LPL, and TNFRSF10B affects plasma fatty acid distribution in Alaskan Eskimos. Voruganti VS, et al. Am | Clin Nutr, 2010 Jun. PMID 20410100. Decreased infiltration of macrophage scavenger receptor-positive cells in initial negative biopsy specimens is correlated with positive repeat biopsies of the prostate. Nonomura N, et al. Cancer Sci, 2010 Jun. PMID 20384632.