

NCOA4 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant NCOA4.

Catalog # AT2981a

Specification

NCOA4 Antibody (monoclonal) (M01) - Product Information

Application	WB, IF
Primary Accession	Q13772
Other Accession	BC012736
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2b kappa
Calculated MW	69726

NCOA4 Antibody (monoclonal) (M01) - Additional Information

Gene ID 8031

Other Names

Nuclear receptor coactivator 4, NCoA-4, Androgen receptor coactivator 70 kDa protein, 70 kDa AR-activator, 70 kDa androgen receptor coactivator, Androgen receptor-associated protein of 70 kDa, Ret-activating protein ELE1, NCOA4, ARA70, ELE1, RFG

Target/Specificity

NCOA4 (AAH12736.1, 1 a.a. ~ 575 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

IF~~1:50~200

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

NCOA4 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


NCOA4 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NCOA4 Antibody (monoclonal) (M01) - Images



Immunofluorescence of monoclonal antibody to NCOA4 on HeLa cell. [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (88.99 KDa) .

NCOA4 Antibody (monoclonal) (M01) - Background

This gene encodes an androgen receptor coactivator. The encoded protein interacts with the androgen receptor in a ligand-dependent manner to enhance its transcriptional activity. Chromosomal translocations between this gene and the ret tyrosine kinase gene, also located on chromosome 10, have been associated with papillary thyroid carcinoma. Alternatively spliced transcript variants have been described. Pseudogenes are present on chromosomes 4, 5, 10, and 14.

NCOA4 Antibody (monoclonal) (M01) - References

Absence of BRAF, NRAS, KRAS, HRAS mutations, and RET/PTC gene rearrangements distinguishes dominant nodules in Hashimoto thyroiditis from papillary thyroid carcinomas. Sadow PM, et al.

Endocr Pathol, 2010 Jun. PMID 20012784. The variant rs1867277 in FOXE1 gene confers thyroid cancer susceptibility through the recruitment of USF1/USF2 transcription factors. Landa I, et al. PLoS Genet, 2009 Sep. PMID 19730683. Transcript level modulates the inherent oncogenicity of RET/PTC oncoproteins. Richardson DS, et al. Cancer Res, 2009 Jun 1. PMID 19487296. Multiple genetic variants along candidate pathways influence plasma high-density lipoprotein cholesterol concentrations. Lu Y, et al. J Lipid Res, 2008 Dec. PMID 18660489. Stimulation of prostate cancer cellular proliferation and invasion by the androgen receptor co-activator ARA70. Peng Y, et al. Am J Pathol, 2008 Jan. PMID 18156210.