

NDN Antibody (monoclonal) (M02)**Mouse monoclonal antibody raised against a partial recombinant NDN.****Catalog # AT2988a****Specification****NDN Antibody (monoclonal) (M02) - Product Information**

Application	WB, IHC, IF, E
Primary Accession	Q99608
Other Accession	NM_002487
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	36086

NDN Antibody (monoclonal) (M02) - Additional Information**Gene ID** 4692**Other Names**

Necdin, NDN

Target/Specificity

NDN (NP_002478, 222 a.a. ~ 321 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

DilutionWB~~1:500~1000
IHC~~1:100~500
IF~~1:50~200
E~~N/A**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions


NDN Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.

NDN Antibody (monoclonal) (M02) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)


- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NDN Antibody (monoclonal) (M02) - Images


Immunofluorescence of monoclonal antibody to NDN on HeLa cell. [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.11 KDa) .




NDN monoclonal antibody (M02), clone 1B3 Western Blot analysis of NDN expression in HL-60 (Cat # AT2988a)




Western Blot analysis of NDN expression in transfected 293T cell line by NDN monoclonal antibody (M02), clone 1B3.

Lane 1: NDN transfected lysate(36.1 KDa).


Lane 2: Non-transfected lysate.



Western blot analysis of NDN over-expressed 293 cell line, cotransfected with NDN Validated Chimera RNAi ((Cat # AT2988a)



Immunoperoxidase of monoclonal antibody to NDN on formalin-fixed paraffin-embedded human colon. [antibody concentration 1.5 ug/ml]



Detection limit for recombinant GST tagged NDN is approximately 0.03ng/ml as a capture antibody.

NDN Antibody (monoclonal) (M02) - Background

This intronless gene is located in the Prader-Willi syndrome deletion region. It is an imprinted gene and is expressed exclusively from the paternal allele. Studies in mouse suggest that the protein encoded by this gene may suppress growth in postmitotic neurons.

NDN Antibody (monoclonal) (M02) - References

- 1.Necdin Enhances Myoblasts Survival by Facilitating the Degradation of the Mediator of Apoptosis CCAR1/CARP1.Francois S, D'Orlando C, Fatone T, Touvier T, Pessina P, Meneveri R, Brunelli S.PLoS One. 2012;7(8):e43335. Epub 2012 Aug 14.