

NFYC Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant NFYC. Catalog # AT3043a

Specification

NFYC Antibody (monoclonal) (M01) - Product Information

Application WB, IHC, IF, E **Primary Accession** 013952 BC005003 Other Accession Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG3 Kappa Calculated MW 50302

NFYC Antibody (monoclonal) (M01) - Additional Information

Gene ID 4802

Other Names

Nuclear transcription factor Y subunit gamma, CAAT box DNA-binding protein subunit C, Nuclear transcription factor Y subunit C, NF-YC, Transactivator HSM-1/2, NFYC

Target/Specificity

NFYC (AAH05003, 14 a.a. \sim 113 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000 IHC~~1:100~500 IF~~1:50~200 E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

NFYC Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

NFYC Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.




- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

NFYC Antibody (monoclonal) (M01) - Images




Immunofluorescence of monoclonal antibody to NFYC on HeLa cell . [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .







Western Blot analysis of NFYC expression in transfected 293T cell line by NFYC monoclonal antibody (M01), clone 1D3.

Lane 1: NFYC transfected lysate(37.2 KDa).

Lane 2: Non-transfected lysate.




Western blot analysis of NFYC over-expressed 293 cell line, cotransfected with NFYC Validated Chimera RNAi ((Cat # AT3043a)



Immunoperoxidase of monoclonal antibody to NFYC on formalin-fixed paraffin-embedded human testis. [antibody concentration 3 ug/ml]





Detection limit for recombinant GST tagged NFYC is 3 ng/ml as a capture antibody.

NFYC Antibody (monoclonal) (M01) - Background

This gene encodes one subunit of a trimeric complex forming a highly conserved transcription factor that binds with high specificity to CCAAT motifs in the promoters of a variety of genes. The encoded protein, subunit C, forms a tight dimer with the B subunit, a prerequisite for subunit A association. The resulting trimer binds to DNA with high specificity and affinity. Subunits B and C each contain a histone-like motif. Multiple transcript variants encoding different isoforms have been found for this gene.

NFYC Antibody (monoclonal) (M01) - References

Replication of previous genome-wide association studies of bone mineral density in premenopausal American women. Ichikawa S, et al. J Bone Miner Res, 2010 Aug. PMID 20200978.NF-YC functions as a corepressor of agonist-bound mineralocorticoid receptor. Murai-Takeda A, et al. J Biol Chem, 2010 Mar 12. PMID 20054001.NF-YC complexity is generated by dual promoters and alternative splicing. Ceribelli M, et al. J Biol Chem, 2009 Dec 4. PMID 19690168.Expression levels of NF-Y target genes changed by CDKN1B correlate with clinical prognosis in multiple cancers. Yamanaka K, et al. Genomics, 2009 Oct. PMID 19559782.Single nucleotide polymorphisms of microRNA machinery genes modify the risk of renal cell carcinoma. Horikawa Y, et al. Clin Cancer Res, 2008 Dec 1. PMID 19047128.