

P2RX5 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant P2RX5.****Catalog # AT3167a****Specification**

P2RX5 Antibody (monoclonal) (M01) - Product Information

Application	WB, E
Primary Accession	O93086
Other Accession	NM_002561
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	49303

P2RX5 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 5026**Other Names**

P2X purinoceptor 5, P2X5, ATP receptor, Purinergic receptor, P2RX5, P2X5

Target/Specificity

P2RX5 (NP_002552, 126 a.a. ~ 224 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

P2RX5 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


P2RX5 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

P2RX5 Antibody (monoclonal) (M01) - Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.63 KDa) .




Western Blot analysis of P2RX5 expression in transfected 293T cell line by P2RX5 monoclonal antibody (M01), clone 1C5.

Lane 1: P2RX5 transfected lysate (46.42 KDa).

Lane 2: Non-transfected lysate.



Western blot analysis of P2RX5 over-expressed 293 cell line, cotransfected with P2RX5 Validated Chimera RNAi (Cat # AT3167a)



Detection limit for recombinant GST tagged P2RX5 is approximately 1ng/ml as a capture antibody.

P2RX5 Antibody (monoclonal) (M01) - Background

The product of this gene belongs to the family of purinoceptors for ATP. This receptor functions as a ligand-gated ion channel. Several characteristic motifs of ATP-gated channels are present in its primary structure, but, unlike other members of the purinoceptors family, this receptor has only a single transmembrane domain. Three transcript variants encoding distinct isoforms have been identified for this gene.

P2RX5 Antibody (monoclonal) (M01) - References

Genetic and functional analysis of human P2X5 reveals a distinct pattern of exon 10 polymorphism with predominant expression of the nonfunctional receptor isoform. Kotnis S, et al. Mol Pharmacol, 2010 Jun. PMID 20223879. Aberrant expression of the hematopoietic-restricted minor histocompatibility antigen LHR-1 on solid tumors results in efficient cytotoxic T cell-mediated lysis. Overes IM, et al. Cancer Immunol Immunother, 2009 Mar. PMID 18719914. Go it alone no more--P2X7 joins the society of heteromeric ATP-gated receptor channels. Dubyak GR. Mol Pharmacol, 2007 Dec. PMID 17895406. P2X5 subunit assembly requires scaffolding by the second transmembrane domain and a conserved aspartate. Duckwitz W, et al. J Biol Chem, 2006 Dec 22. PMID 17001079. Purinoceptor expression on keratinocytes reflects their function on the epidermis during chronic venous insufficiency. Metcalfe MJ, et al. Arch Dermatol Res, 2006 Nov. PMID 16967306.