

PARC Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant PARC.****Catalog # AT3180a****Specification**

PARC Antibody (monoclonal) (M01) - Product Information

Application	WB
Primary Accession	Q8IWT3
Other Accession	NM_015089
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	281229

PARC Antibody (monoclonal) (M01) - Additional Information**Gene ID** 23113**Other Names**

Cullin-9, CUL-9, Ubch7-associated protein 1, p53-associated parkin-like cytoplasmic protein, CUL9, H7AP1, KIAA0708, PARC

Target/Specificity

PARC (NP_055904, 918 a.a. ~ 1025 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

PARC Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


PARC Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PARC Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.62 kDa) .

PARC Antibody (monoclonal) (M01) - References

Association of genetic variants with hemorrhagic stroke in Japanese individuals. Yoshida T, et al. Int J Mol Med, 2010 Apr. PMID 20198315. Assessment of a polymorphism of SDK1 with hypertension in Japanese Individuals. Oguri M, et al. Am J Hypertens, 2010 Jan. PMID 19851296. Activation of p53-dependent responses in tumor cells treated with a PARC-interacting peptide. Vitali R, et al. Biochem Biophys Res Commun, 2008 Apr 4. PMID 18230339. Systematic identification of SH3 domain-mediated human protein-protein interactions by peptide array target screening. Wu C, et al. Proteomics, 2007 Jun. PMID 17474147. Hyperubiquitylation of wild-type p53 contributes to cytoplasmic sequestration in neuroblastoma. Becker K, et al. Cell Death Differ, 2007 Jul. PMID 17380154.