

PBK Antibody (monoclonal) (M06)
Mouse monoclonal antibody raised against a partial recombinant PBK.
Catalog # AT3196a

Specification

PBK Antibody (monoclonal) (M06) - Product Information

Application	WB, IHC, IF
Primary Accession	Q96KB5
Other Accession	BC015191
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2b Kappa
Calculated MW	36085

PBK Antibody (monoclonal) (M06) - Additional Information

Gene ID 55872

Other Names

Lymphokine-activated killer T-cell-originated protein kinase, Cancer/testis antigen 84, CT84, MAPKK-like protein kinase, Nori-3, PDZ-binding kinase, Spermatogenesis-related protein kinase, SPK, T-LAK cell-originated protein kinase, PBK, TOPK

Target/Specificity

PBK (AAH15191, 1 a.a. ~ 110 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

IHC~~1:100~500

IF~~1:50~200

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions


PBK Antibody (monoclonal) (M06) is for research use only and not for use in diagnostic or therapeutic procedures.

PBK Antibody (monoclonal) (M06) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PBK Antibody (monoclonal) (M06) - Images




Immunofluorescence of monoclonal antibody to PBK on HeLa cell. [antibody concentration 10 ug/ml]



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.73 KDa) .



PBK monoclonal antibody (M06), clone 4A10 Western Blot analysis of PBK expression in HeLa S3 NE (Cat # AT3196a)



Immunoperoxidase of monoclonal antibody to PBK on formalin-fixed paraffin-embedded human testis. [antibody concentration 3 ug/ml]

PBK Antibody (monoclonal) (M06) - Background

This gene encodes a serine/threonine kinase related to the dual specific mitogen-activated protein kinase kinase (MAPKK) family. Evidence suggests that mitotic phosphorylation is required for its catalytic activity. This mitotic kinase may be involved in the activation of lymphoid cells and support testicular functions, with a suggested role in the process of spermatogenesis.

PBK Antibody (monoclonal) (M06) - References

Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. PBK/TOPK in the differential diagnosis of cholangiocarcinoma from hepatocellular carcinoma and its involvement in prognosis of human cholangiocarcinoma. He F, et al. Hum Pathol, 2010 Mar. PMID 19954816. Critical roles of T-LAK cell-originated protein kinase in cytokinesis. Park JH, et al. Cancer Sci, 2010 Feb. PMID 19900192. Molecular genetics of successful smoking cessation: convergent genome-wide association study results. Uhl GR, et al. Arch Gen Psychiatry, 2008 Jun. PMID 18519826. Bidirectional signals transduced by TOPK-ERK interaction increase tumorigenesis of HCT116 colorectal cancer cells. Zhu F, et al. Gastroenterology, 2007 Jul. PMID 17631144.