

PCDHA10 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant PCDHA10.****Catalog # AT3217a****Specification**

PCDHA10 Antibody (monoclonal) (M01) - Product Information

Application	WB, IHC
Primary Accession	O9Y5I2
Other Accession	NM_018901
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	102875

PCDHA10 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 56139**Other Names**

Protocadherin alpha-10, PCDH-alpha-10, PCDHA10, CNRS8

Target/Specificity

PCDHA10 (NP_061724, 182 a.a. ~ 290 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

IHC~~1:100~500

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

PCDHA10 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


PCDHA10 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PCDHA10 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.73 kDa) .



Immunoperoxidase of monoclonal antibody to PCDHA10 on formalin-fixed paraffin-embedded human stomach. [antibody concentration 3 ug/ml]

PCDHA10 Antibody (monoclonal) (M01) - Background

This gene is a member of the protocadherin alpha gene cluster, one of three related gene clusters tandemly linked on chromosome five that demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The alpha gene cluster is composed of 15 cadherin superfamily genes related to the mouse CNR genes and consists of 13 highly similar and 2 more distantly related coding sequences. The tandem array of 15 N-terminal exons, or variable exons, are followed by downstream C-terminal exons, or constant exons, which are shared by all genes in the cluster. The large, uninterrupted N-terminal exons each encode six cadherin ectodomains while the C-terminal exons encode the cytoplasmic domain. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins that most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been observed and additional variants have been suggested but their full-length nature has yet to be determined.

PCDHA10 Antibody (monoclonal) (M01) - References

Systematic identification of SH3 domain-mediated human protein-protein interactions by peptide array target screening. Wu C, et al. Proteomics, 2007 Jun. PMID 17474147. The DNA sequence and comparative analysis of human chromosome 5. Schmutz J, et al. Nature, 2004 Sep 16. PMID 15372022. Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932. Comparative DNA sequence analysis of mouse and human protocadherin gene clusters. Wu Q, et al. Genome Res, 2001 Mar. PMID 11230163. Phylogenetic analysis of the cadherin superfamily allows identification of six major subfamilies besides several solitary members. Nollet F, et al. J Mol Biol, 2000 Jun 9. PMID 10835267.