

**PCDHB16 Antibody (monoclonal) (M03)****Mouse monoclonal antibody raised against a full length recombinant PCDHB16.****Catalog # AT3224a****Specification**

---

**PCDHB16 Antibody (monoclonal) (M03) - Product Information**

Application	WB
Primary Accession	<a href="#">Q9NRJ7</a>
Other Accession	<a href="#">BC036062</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2b Kappa
Calculated MW	84936

**PCDHB16 Antibody (monoclonal) (M03) - Additional Information****Gene ID** 57717**Other Names**

Protocadherin beta-16, PCDH-beta-16, Protocadherin-3X, PCDHB16, KIAA1621, PCDH3X

**Target/Specificity**

PCDHB16 (AAH36062.1, 1 a.a. ~ 776 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

PCDHB16 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.


**PCDHB16 Antibody (monoclonal) (M03) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### PCDHB16 Antibody (monoclonal) (M03) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (111.1 kDa) .

### PCDHB16 Antibody (monoclonal) (M03) - Background

This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the transcripts from these genes are made up of only one large exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment and function of specific cell-cell neural connections.

### PCDHB16 Antibody (monoclonal) (M03) - References

The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334. Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. Nat Genet, 2004 Jan. PMID 14702039. Exploring proteomes and analyzing protein processing by mass spectrometric identification of sorted N-terminal peptides. Gevaert K, et al. Nat Biotechnol, 2003 May. PMID 12665801. Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932. Protocadherins. Frank M, et al. Curr Opin Cell Biol, 2002 Oct. PMID 12231349.