

PCDHB3 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant PCDHB3.

Catalog # AT3225a

Specification

PCDHB3 Antibody (monoclonal) (M01) - Product Information

Application	WB, E
Primary Accession	O9Y5E6
Other Accession	NM_018937
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	86714

PCDHB3 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 56132**Other Names**

Protocadherin beta-3, PCDH-beta-3, PCDHB3

Target/Specificity

PCDHB3 (NP_061760, 284 a.a. ~ 381 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB ~ 1:500 ~ 1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

PCDHB3 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


PCDHB3 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PCDHB3 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.52 KDa) .



Detection limit for recombinant GST tagged PCDHB3 is approximately 0.3ng/ml as a capture antibody.

PCDHB3 Antibody (monoclonal) (M01) - Background

This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the transcripts from these genes are made up of only one large exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment and function of specific cell-cell neural connections.

PCDHB3 Antibody (monoclonal) (M01) - References

Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932. Protocadherins. Frank M, et al. Curr Opin Cell Biol, 2002 Oct. PMID 12231349. The human and murine protocadherin-beta one-exon gene families show high evolutionary conservation, despite the difference in gene

number. Vanhalst K, et al. FEBS Lett, 2001 Apr 20. PMID 11322959. Comparative DNA sequence analysis of mouse and human protocadherin gene clusters. Wu Q, et al. Genome Res, 2001 Mar. PMID 11230163. Phylogenetic analysis of the cadherin superfamily allows identification of six major subfamilies besides several solitary members. Nollet F, et al. J Mol Biol, 2000 Jun 9. PMID 10835267.