

**PIGO Antibody (monoclonal) (M01)****Mouse monoclonal antibody raised against a full length recombinant PIGO.****Catalog # AT3307a****Specification**

---

**PIGO Antibody (monoclonal) (M01) - Product Information**

Application	WB, E
Primary Accession	<a href="#">Q8TEQ8</a>
Other Accession	<a href="#">BC001030</a>
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2b kappa
Calculated MW	118699

**PIGO Antibody (monoclonal) (M01) - Additional Information****Gene ID** 84720**Other Names**

GPI ethanolamine phosphate transferase 3, 2---, Phosphatidylinositol-glycan biosynthesis class O protein, PIG-O, PIGO

**Target/Specificity**

PIGO (AAH01030, 1 a.a. ~ 454 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

**Dilution**

WB~~1:500~1000

**Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

**Storage**

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Precautions**

PIGO Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


**PIGO Antibody (monoclonal) (M01) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### PIGO Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (75.68 kDa) .



Detection limit for recombinant GST tagged PIGO is approximately 0.1 ng/ml as a capture antibody.

### PIGO Antibody (monoclonal) (M01) - Background

This gene encodes a protein that is involved in glycosylphosphatidylinositol (GPI)-anchor biosynthesis. The GPI-anchor is a glycolipid which contains three mannose molecules in its core backbone. The GPI-anchor is found on many blood cells and serves to anchor proteins to the cell surface. This protein is involved in the transfer of ethanolaminephosphate (EtNP) to the third mannose in GPI. At least two alternatively spliced transcripts encoding distinct isoforms have been found for this gene.

### PIGO Antibody (monoclonal) (M01) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolidinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614. Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC).

Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334. DNA sequence and analysis of human chromosome 9. Humphray SJ, et al. Nature, 2004 May 27. PMID 15164053.