

PIGO Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a full length recombinant PIGO. Catalog # AT3307a

Specification

PIGO Antibody (monoclonal) (M01) - Product Information

Application WB, E **Primary Accession Q8TEQ8** Other Accession BC001030 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG2b kappa Calculated MW 118699

PIGO Antibody (monoclonal) (M01) - Additional Information

Gene ID 84720

Other Names

GPI ethanolamine phosphate transferase 3, 2---, Phosphatidylinositol-glycan biosynthesis class O protein, PIG-O, PIGO

Target/Specificity

PIGO (AAH01030, 1 a.a. \sim 454 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000 E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

PIGO Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

PIGO Antibody (monoclonal) (M01) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides




- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

PIGO Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (75.68 KDa).



Detection limit for recombinant GST tagged PIGO is approximately 0.1ng/ml as a capture antibody.

PIGO Antibody (monoclonal) (M01) - Background

This gene encodes a protein that is involved in glycosylphosphatidylinositol (GPI)-anchor biosynthesis. The GPI-anchor is a glycolipid which contains three mannose molecules in its core backbone. The GPI-anchor is found on many blood cells and serves to anchor proteins to the cell surface. This protein is involved in the transfer of ethanolaminephosphate (EtNP) to the third mannose in GPI. At least two alternatively spliced transcripts encoding distinct isoforms have been found for this gene.

PIGO Antibody (monoclonal) (M01) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121.The status,







quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.DNA sequence and analysis of human chromosome 9. Humphray SJ, et al. Nature, 2004 May 27. PMID 15164053.