

PLCD4 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant PLCD4.****Catalog # AT3334a****Specification**

PLCD4 Antibody (monoclonal) (M01) - Product Information

Application	WB, E
Primary Accession	Q9BRC7
Other Accession	BC006355
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 Kappa
Calculated MW	87585

PLCD4 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 84812**Other Names**

1-phosphatidylinositol 4, 5-bisphosphate phosphodiesterase delta-4, hPLCD4, Phosphoinositide phospholipase C-delta-4, Phospholipase C-delta-4, PLC-delta-4, PLCD4

Target/Specificity

PLCD4 (AAH06355, 18 a.a. ~ 127 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

PLCD4 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


PLCD4 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PLCD4 Antibody (monoclonal) (M01) - Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.84 kDa) .



Western Blot analysis of PLCD4 expression in transfected 293T cell line by PLCD4 monoclonal antibody (M01), clone 4D4.

Lane 1: PLCD4 transfected lysate (Predicted MW: 87.6 kDa).

Lane 2: Non-transfected lysate.



Detection limit for recombinant GST tagged PLCD4 is approximately 0.3ng/ml as a capture antibody.

PLCD4 Antibody (monoclonal) (M01) - Background

Phosphatidylinositol-specific phospholipase C (PLC) plays an important role in receptor-mediated signal transduction by generating 2 second messenger molecules, inositol 1,4,5-triphosphate (IP3) and diacylglycerol, from phosphatidylinositol 4,5-bisphosphate (PIP2). PLC comprises a diverse family of enzymes that differ in structure and tissue distribution (Berridge, 1993 [PubMed 8381210]).

PLCD4 Antibody (monoclonal) (M01) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086. Polymorphisms in innate immunity genes and patients response to dendritic cell-based HIV immuno-treatment. Segat L, et al. Vaccine, 2010 Mar 2. PMID 20056178. Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121. Identification of new putative susceptibility genes for several psychiatric disorders by association analysis of regulatory and non-synonymous SNPs of 306 genes involved in neurotransmission and neurodevelopment. Gratac?s M, et al. Am J Med Genet B Neuropsychiatr Genet, 2009 Sep 5. PMID 19086053. Identification and analysis of the promoter region of the human PLC-delta4 gene. Chae SW, et al. Mol Biol Rep, 2007 Jun. PMID 17394098.