

RASA1 Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant RASA1.****Catalog # AT3574a****Specification**

RASA1 Antibody (monoclonal) (M01) - Product Information

Application	WB, IHC, E
Primary Accession	P20936
Other Accession	NM_002890
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG1 kappa
Calculated MW	116403

RASA1 Antibody (monoclonal) (M01) - Additional Information**Gene ID** 5921**Other Names**

Ras GTPase-activating protein 1, GAP, GTPase-activating protein, RasGAP, Ras p21 protein activator, p120GAP, RASA1, GAP, RASA

Target/Specificity

RASA1 (NP_002881, 948 a.a. ~ 1047 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

IHC~~1:100~500

E~~N/A

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

RASA1 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


RASA1 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


RASA1 Antibody (monoclonal) (M01) - Images




Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .



RASA1 monoclonal antibody (M01), clone 2C12 Western Blot analysis of RASA1 expression in IMR-32 ((Cat # AT3574a)



Immunoperoxidase of monoclonal antibody to RASA1 on formalin-fixed paraffin-embedded human small Intestine. [antibody concentration 6 ug/ml]



Detection limit for recombinant GST tagged RASA1 is 0.3 ng/ml as a capture antibody.

RASA1 Antibody (monoclonal) (M01) - Background

The protein encoded by this gene is located in the cytoplasm and is part of the GAP1 family of GTPase-activating proteins. The gene product stimulates the GTPase activity of normal RAS p21 but not its oncogenic counterpart. Acting as a suppressor of RAS function, the protein enhances the weak intrinsic GTPase activity of RAS proteins resulting in the inactive GDP-bound form of RAS, thereby allowing control of cellular proliferation and differentiation. Mutations leading to changes in the binding sites of either protein are associated with basal cell carcinomas. Alternative splicing results in two isoforms where the shorter isoform, lacking the N-terminal hydrophobic region but retaining the same activity, appears to be abundantly expressed in placental but not adult tissues.

RASA1 Antibody (monoclonal) (M01) - References

Identification of several novel non-p.R132 IDH1 variants in thyroid carcinomas. Hemery JP, et al. Eur J Endocrinol, 2010 Aug 11. PMID 20702649. Backtracking RAS mutations in high hyperdiploid childhood acute lymphoblastic leukemia. Wiemels JL, et al. Blood Cells Mol Dis, 2010 Oct 15. PMID 20688547. MicroRNA-132-mediated loss of p120RasGAP activates the endothelium to facilitate pathological angiogenesis. Anand S, et al. Nat Med, 2010 Aug. PMID 20676106. High frequency type I/II mutational shifts between diagnosis and relapse are associated with outcome in pediatric AML: implications for personalized medicine. Bachas C, et al. Blood, 2010 Jun 30. PMID 20592250. A novel association between RASA1 mutations and spinal arteriovenous anomalies. Thiex R, et al. AJNR Am J Neuroradiol, 2010 Apr. PMID 20007727.