

# RASA3 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant RASA3. Catalog # AT3575a

# **Specification**

# RASA3 Antibody (monoclonal) (M01) - Product Information

**Application** WB, E **Primary Accession** 014644 Other Accession BC047242 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG2a kappa Calculated MW 95699

# RASA3 Antibody (monoclonal) (M01) - Additional Information

#### **Gene ID 22821**

### **Other Names**

Ras GTPase-activating protein 3, GAP1(IP4BP), Ins P4-binding protein, RASA3

## **Target/Specificity**

RASA3 (AAH47242, 725 a.a. ~ 834 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

# Dilution

WB~~1:500~1000

E~~N/A

## **Format**

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

#### **Precautions**

RASA3 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

### RASA3 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot




- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# RASA3 Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (37.73 KDa).




Western Blot analysis of RASA3 expression in transfected 293T cell line by RASA3 monoclonal antibody (M01), clone 1F11.

Lane 1: RASA3 transfected lysate (Predicted MW: 95.7 KDa).

Lane 2: Non-transfected lysate.





Detection limit for recombinant GST tagged RASA3 is approximately 0.3ng/ml as a capture antibody.

# RASA3 Antibody (monoclonal) (M01) - Background

The protein encoded by this gene is member of the GAP1 family of GTPase-activating proteins. The gene product stimulates the GTPase activity of normal RAS p21 but not its oncogenic counterpart. Acting as a suppressor of RAS function, the protein enhances the weak intrinsic GTPase activity of RAS proteins resulting in the inactive GDP-bound form of RAS, thereby allowing control of cellular proliferation and differentiation. This family member is an inositol 1,3,4,5-tetrakisphosphate-binding protein, like the closely related RAS p21 protein activator 2. The two family members have distinct pleckstrin-homology domains, with this particular member having a domain consistent with its localization to the plasma membrane.

# RASA3 Antibody (monoclonal) (M01) - References

The ability of GAP1IP4BP to function as a Rap1 GTPase-activating protein (GAP) requires its Ras GAP-related domain and an arginine finger rather than an asparagine thumb. Kupzig S, et al. Mol Cell Biol, 2009 Jul. PMID 19433443.Immunoaffinity profiling of tyrosine phosphorylation in cancer cells. Rush J, et al. Nat Biotechnol, 2005 Jan. PMID 15592455.Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. Nat Genet, 2004 Jan. PMID 14702039.Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932.Analyzing the role of the putative inositol 1,3,4,5-tetrakisphosphate receptor GAP1IP4BP in intracellular Ca2+homeostasis. Walker SA, et al. J Biol Chem, 2002 Dec 13. PMID 12356770.