

RBAK Antibody (monoclonal) (M01)**Mouse monoclonal antibody raised against a partial recombinant RBAK.****Catalog # AT3585a****Specification**

RBAK Antibody (monoclonal) (M01) - Product Information

Application	WB, E
Primary Accession	O9NYW8
Other Accession	NM_021163
Reactivity	Human
Host	mouse
Clonality	Monoclonal
Isotype	IgG2a Kappa
Calculated MW	82995

RBAK Antibody (monoclonal) (M01) - Additional Information**Gene ID** 57786**Other Names**

RB-associated KRAB zinc finger protein, RB-associated KRAB repressor, hRBaK, Zinc finger protein 769, RBAK, ZNF769

Target/Specificity

RBAK (NP_066986, 51 a.a. ~ 150 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

RBAK Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.


RBAK Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RBAK Antibody (monoclonal) (M01) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 kDa) .



Detection limit for recombinant GST tagged RBAK is approximately 0.03ng/ml as a capture antibody.

RBAK Antibody (monoclonal) (M01) - Background

This gene encodes a nuclear protein which interacts with the tumor suppressor retinoblastoma 1. The two interacting proteins are thought to act as a transcriptional repressor for promoters which are activated by the E2F1 transcription factor. This protein contains a Kruppel-associated box (KRAB), which is a transcriptional repressor motif.

RBAK Antibody (monoclonal) (M01) - References

Identification of candidate genes for human pituitary development by EST analysis. Ma Y, et al. BMC Genomics, 2009 Mar 15. PMID 19284880. Coeliac disease-associated risk variants in TNFAIP3 and REL implicate altered NF-kappaB signalling. Trynka G, et al. Gut, 2009 Aug. PMID 19240061. Examination of chromosome 7p22 candidate genes RBAK, PMS2 and GNA12 in familial hyperaldosteronism type II. Jeske YW, et al. Clin Exp Pharmacol Physiol, 2008 Apr. PMID 18307725. No evidence for coding region mutations in the retinoblastoma-associated Kruppel-associated box protein gene (RBAK) causing familial hyperaldosteronism type II. So A, et al. Clin Endocrinol (Oxf), 2006 Dec. PMID 17121540. Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes.

Kimura K, et al. Genome Res, 2006 Jan. PMID 16344560.