

RFP2 Antibody (monoclonal) (M01)

Mouse monoclonal antibody raised against a partial recombinant RFP2. Catalog # AT3624a

Specification

RFP2 Antibody (monoclonal) (M01) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW E <u>Q8N513</u> <u>NM_005798</u> Human mouse Monoclonal IgG2a lambda 31048

RFP2 Antibody (monoclonal) (M01) - Additional Information

Gene ID 283518

Other Names Potassium channel regulatory protein, Potassium channel regulator, Protein CLLD4, KCNRG, CLLD4

Target/Specificity RFP2 (NP_005789, 1 a.a. ~ 100 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution E~~N/A

Format Clear, colorless solution in phosphate buffered saline, pH 7.2 .

Storage Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions RFP2 Antibody (monoclonal) (M01) is for research use only and not for use in diagnostic or therapeutic procedures.

RFP2 Antibody (monoclonal) (M01) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry



Immunofluorescence

- <u>Immunoprecipitation</u>
- Flow Cytomety
- <u>Cell Culture</u>

RFP2 Antibody (monoclonal) (M01) - Images



Detection limit for recombinant GST tagged RFP2 is approximately 0.1ng/ml as a capture antibody.

RFP2 Antibody (monoclonal) (M01) - Background

This gene encodes a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This gene is located on chromosome 13 within the minimal deletion region for B-cell chronic lymphocytic leukemia. Multiple alternatively spliced transcript variants have been found for this gene.

RFP2 Antibody (monoclonal) (M01) - References

Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348. The RBCC gene RFP2 (Leu5) encodes a novel transmembrane E3 ubiquitin ligase involved in ERAD. Lerner M, et al. Mol Biol Cell, 2007 May. PMID 17314412. Human RFP2 gene promoter: unique structure and unusual strength. Skoblov M, et al. Biochem Biophys Res Commun, 2006 Apr 14. PMID 16499869. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334. DLEU2 encodes an antisense RNA for the putative bicistronic RFP2/LEU5 gene in humans and mouse. Corcoran MM, et al. Genes Chromosomes Cancer, 2004 Aug. PMID 15188451.